



**WHO/CDS/CSR/ISR/2001.2**

**Protocol for the Assessment of National Communicable  
Disease Surveillance and Response Systems**

**Guidelines for Assessment Teams**

**World Health Organization**  
Department of Communicable Disease  
Surveillance and Response

This document has been downloaded from the WHO/EMC Web site. The original cover pages and lists of participants are not included. See <http://www.who.int/emc> for more information.

**© World Health Organization**

This document is not a formal publication of the World Health Organization (WHO), and all rights are reserved by the Organization. The document may, however, be freely reviewed, abstracted, reproduced and translated, in part or in whole, but not for sale nor for use in conjunction with commercial purposes.

The views expressed in documents by named authors are solely the responsibility of those authors. The mention of specific companies or specific manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned.

# TABLE OF CONTENTS

---

|   |     |
|---|-----|
| TABLE OF CONTENTS .....   | i   |
| ACKNOWLEDGEMENTS.....   | iii |
| ACRONYMS .....  | v   |
| INTRODUCTION.....   | 1   |
| WHAT SHOULD THIS DOCUMENT BE USED FOR? .....  | 1   |
| WHAT IS THE NATIONAL SURVEILLANCE SYSTEM?.....  | 1   |
| WHY ASSESS THE NATIONAL SURVEILLANCE SYSTEM?.....   | 2   |
| WHAT IS A MULTI-DISEASE OR AN INTEGRATED APPROACH TO<br>DISEASE SURVEILLANCE? .....             | 2   |
| WHAT ARE THE AIMS AND OBJECTIVES OF THE ASSESSMENT? .....                                       | 4   |
| WHAT SHOULD BE ASSESSED? .....  | 4   |
| WHAT SHOULD GUIDE THE ASSESSMENT? .....   | 8   |
| PROCEDURES, ACTIVITIES AND TIMETABLE OF THE ASSESSMENT.....                                     | 8   |
| PHASE I PLANNING THE MISSION .....  | 9   |
| PHASE II THE ASSESSMENT .....   | 11  |
| STEP 1 PRE-ASSESSMENT FACILITATED WORKSHOP<br>WITH NATIONAL TEAM.....                           | 11  |
| STEP 2 TRAINING OF ASSESSMENT TEAMS.....  | 12  |
| STEP 3 FIELD ASSESSMENT .....   | 12  |
| STEP 4 ANALYSIS AND PRELIMINARY REPORT WRITING .....  | 13  |
| STEP 5 POST-ASSESSMENT WORKSHOP TO PRESENT<br>PRELIMINARY FINDINGS.....                         | 15  |
| PHASE III WORKSHOP TO ELABORATE PLAN OF ACTION .....  | 16  |
| PHASE IV FOLLOW-UP OF THE IMPLEMENTATION OF THE<br>MULTI-DISEASE APPROACH TO SURVEILLANCE ..... | 18  |
| LIST OF ANNEXES .....   | 19  |



## ACKNOWLEDGEMENTS

---

*The World Health Organization wishes to acknowledge the support of the United States Agency for International Development, the United Nations Foundation for International Partnerships, the Department for International Development of the United Kingdom and the Government of Ireland in the production of this document.*

*The World Health Organization is also grateful for technical support in the completion of this work from the WHO Regional Office for Africa and the Capacity Development Branch, Division of International Health, Epidemiology Program Office, Centers for Disease Control and Prevention, United States.*

*We also thank the other WHO Regional Offices, particularly EMRO, EURO and SEARO where the protocol has been field tested.*



## **ACRONYMS**

---

|                 |   |
|-----------------|---|
| <b>ADB</b>      | African Development Bank                                  |
| <b>AFP</b>      | Acute Flaccid Paralysis                                   |
| <b>AIDS</b>     | Acquired Immune Deficiency Syndrome                       |
| <b>CDC</b>      | Centers for Disease Control and Prevention                |
| <b>DANIDA</b>   | Danish Development Aid                                    |
| <b>DFID</b>     | Department for International Development                  |
| <b>EU</b>       | European Union  |
| <b>FETP</b>     | Field Epidemiology Training Programme                     |
| <b>GIS</b>      | Geographic Information System                             |
| <b>HIV</b>      | Human Immunodeficiency Virus                              |
| <b>HQ</b>       | World Health Organization Headquarters                    |
| <b>IDS</b>      | Integrated Disease Surveillance                           |
| <b>MoH</b>      | Ministry of Health  |
| <b>NGO</b>      | Non-Governmental Organization                             |
| <b>PoA</b>      | Plan of Action  |
| <b>RO</b>       | World Health Organization Regional Office                 |
| <b>TB</b>       | Tuberculosis  |
| <b>UN</b>       | United Nations  |
| <b>UNAIDS</b>   | United Nations Programme on HIV/AIDS                      |
| <b>UNDP</b>     | United Nations Development Programme                      |
| <b>UNICEF</b>   | United Nations Children's Fund                            |
| <b>USAID</b>    | United States Agency for International Development        |
| <b>WHO</b>      | World Health Organization                                 |
| <b>WHO/AFRO</b> | World Health Organization Regional Office for Africa      |
| <b>WR</b>       | World Health Organization Country Representative          |
| <b>WRO</b>      | World Health Organization Country Representative's Office |



# INTRODUCTION

---

## **What should this document be used for?**

This manual has been developed for World Health Organization (WHO) staff and partners carrying out assessments of national communicable disease surveillance systems with a national team. It will help WHO staff and consultants guide a group of national professionals through an assessment of the overall structure and performance of surveillance activities in a Member State. This assessment should lead to a standardised report and an agreed plan of action. The plan of action will include a practical timetable for implementation, agreed upon by the Ministry of Health (MoH), WHO and by other partners who may be contributing to the process.

This generic document represents a prototype for the assessment of surveillance and response systems, and may require adaptation in the field. It contains guidance on planning and carrying out an assessment with practical tools such as work group exercises, tables shells and spreadsheets for data collection. The manual also outlines a suggested reporting format with tables for implementation plans.

## **What is the national surveillance system?**

Surveillance is the process of systematic collection, collation and analysis of data with prompt dissemination to those who need to know, for relevant action to be taken. A well functioning disease surveillance system provides information for planning, implementation, monitoring and evaluation of public health intervention programmes. Surveillance for communicable diseases is a part of public health surveillance, which in turn is part of the wider health information system. The objective of the surveillance system and use of the information determines the data collected and the speed of information flow within the system. Early warning of epidemics is essential for effective and rapid control, while information on endemic communicable disease is essential for monitoring the disease. Either way, information on priority communicable diseases is critical for control. Many countries have developed surveillance capacities to monitor diseases with a high burden, to detect outbreaks of epidemic-prone disease and to monitor progress towards national or international control or eradication targets. In this sense, surveillance of communicable diseases is a national function.

### **Why assess the national surveillance system?**

Many countries' surveillance systems have developed in an uneven way, with various surveillance activities funded and managed by different control programmes sometimes based in different institutions (ex. MoH, academic or research institutes, NGOs). Some vertical programmes have kept the surveillance function close to the control function, which is essentially good for the control of a specific disease. On the other hand, overall surveillance functions in a country can become badly disjointed and inefficient. In such cases, field workers participate in multiple systems, use different surveillance methods, terminology, reporting forms and frequency, based on varied training received. This approach may result in extra costs and often leads to work overload and de-motivation for the health worker.

In some cases surveillance is far removed from the control efforts: data are collected on a large number of health events, many of which do not constitute priorities for the country. Detection and reporting of cases and epidemics are rarely carried out on time, and analysis, interpretation and use of available data at all levels for decision making and action is poor.

Each country needs to periodically assess its overall surveillance system so that this continues to reflect national disease control priorities, remains efficient and takes advantages of opportunities for the integration of activities. New surveillance methods and techniques that improve the efficiency of the system should be considered and included in the surveillance system strengthening process.

The World Health Organization (WHO) is promoting a more co-ordinated and synergistic approach to the surveillance and control of communicable diseases. With this in mind, the proposed assessment attempts to deliver an integrated system, using practical and participatory approaches.

### **What is a multi-disease or integrated approach to disease surveillance?**

Surveillance activities for different diseases involve similar functions and very often use the same structures, processes and personnel. A multi-disease approach to disease surveillance aims at establishing well co-ordinated action-oriented surveillance systems that seek opportunities for integration of core and support surveillance functions when appropriate, maximize synergies, take advantage of new tools, build on existing resources, and benefit from successful initiatives. This permits sharing of experiences and resources, avoids duplication of efforts, reduces work load at lower levels, addresses the needs of programmes, and focuses efforts. This approach calls for a co-ordinated approach to data collection, analysis, interpretation and dissemination. It

envisages integration of surveillance activities at all levels when appropriate, while support targeted to surveillance are streamlined and directed in a co-ordinated way.

Disease surveillance should be based on collecting only the information that is required to achieve the control objectives. The data required may differ from disease to disease. For example, the rate of treatment completion and the cure rate are essential indicators in TB surveillance; in HIV/AIDS surveillance the proportion of the population positive for HIV should be monitored as well as the number of new cases of AIDS. Although surveillance may have very specific information needs, many elements of data collection are very similar and the data source is often the same individual or facility. The challenge is to identify where synergy is possible, and exploit this, while at the same time recognizing the needs of some programmes for supplementary information or alternative methods of surveillance.

Specialized surveillance systems (e.g. for acute flaccid paralysis — AFP, or for HIV/AIDS) are important, especially when surveillance methods are complex and the systems have specific information needs. All surveillance systems however, involve the same universal functions (case detection, confirmation, reporting, analysis, investigation, response, feedback and monitoring), and common support functions, (e.g. training, supervision, communications, other resources). It is possible to look at the system as a whole and approach development and strengthening in a co-ordinated way. Opportunities to use common reporting forms, the use of one simple data entry system for multiple diseases and recourse to common communication channels need to be explored. Where possible, all reports should go from district level to a single office at national level. Training and supervision should be integrated and a common feedback bulletin used. Computers, vehicles, fridges etc. can be shared. Instead of competing for funds, different surveillance programmes can work together in appealing for funds.

There may also be differences in the speed at which data and information flow through the system, and the speed of response required for that information. Thus, for the system to function as an “early warning system”, reporting, confirmation, decision making and response should be rapid. On the other hand, for endemic diseases, the aim may be to carefully consider the data collected in order to adjust or target the control programme. The national surveillance system should therefore be able to accommodate both needs, and may require more than one speed for reporting.

In other situations, surveillance that is well developed in one programme may act as a “driving force”, leading to the improvement of other

surveillance activities. It is important to identify these “driving forces” during assessment and to take advantage of them.

### **What are the aims and objectives of the assessment?**

The current approach brings together all those in a country who have responsibility for the surveillance of communicable diseases, with the aim of formally assessing the national disease surveillance systems to strengthen them, using an integrated or multi-disease approach. This assessment should lead to an agreed prioritised plan of action for bringing about improvements in system performance that address gaps identified during the assessment.

The objectives of the assessment are:

1. To obtain baseline information for implementing a co-ordinated, multi-disease approach to disease surveillance that allows measurement of progress made in surveillance strengthening efforts
2. To determine country needs as regards strengthening the surveillance system for communicable disease prevention and control
3. To identify gaps and opportunities in performing the core and support functions of surveillance, and assessing the resources available for these
4. To enable the development of a prioritised action plan, based on the assessment findings.

### **What should be assessed?**

The team should decide on the priority diseases for surveillance and response.

The assessment will be with regard to the structure, organization, processes and output of surveillance and response systems. The capacity for core functions and support functions of surveillance and response at every level of the health care system will be examined. Both core functions and support functions are matched against objectives outlined in a pre-assessment workshop. Opportunities to integrate, co-ordinate and synergize surveillance should be identified during the whole process of assessment, as well as the possibility to use new techniques such as health mapping for surveillance. The attributes of a good surveillance system should be considered (simplicity, flexibility, acceptability, sensitivity, predictive value positive, representativeness, and timeliness) as well as the cost of the system (See Annex 1 for definitions).

### **1. Priority diseases**

Surveillance should ideally centre on priority diseases within the country. Many countries engage in the surveillance of a very large number of diseases. The number of diseases under surveillance continually increases, but the need for this surveillance is often not assessed. In other countries these lists have been inherited from previous administrations. Any assessment of national surveillance should examine all the entities under surveillance and ask the question “is this activity a priority?” Many surveillance systems have a long history where new diseases have been added, while diseases that are no longer a priority have not been deleted. In other cases, countries may lack surveillance in critical areas, especially as diseases can emerge over time as problems that were unforeseen when surveillance was initially developed.

### **2. Assessing structure**

The organization of the surveillance and response systems should be described at the central, intermediate, district, health facility levels and the community level where appropriate. The relationship between the different levels should be described and discussed, as well as the resources (input) that are used for activities at these levels.

### **3. Assessing processes and capacity for surveillance and response**

For each priority disease or group of diseases, the capacity to carry out core and support functions of surveillance and response should be reviewed. The procedure for information flow should be described and its use for public health action assessed. Duplication in the implementation of these functions should be noted. The capacity of the national surveillance system is determined by the ability of the system to monitor priority health events adequately.

The core activities and support functions of the surveillance system will be assessed at all levels of health care (central, regional/provincial, district or equivalent, health facility). The core activities for an effective surveillance for any health event are:

- Detection (identifying cases and outbreaks)
- Registration
- Confirmation (epidemiological and laboratory confirmation)
- Reporting (early warning and routine)
- Analysis and interpretation (preparing and periodically updating graphs, tables and charts to describe time, person and place for reported diseases and conditions, identifying unusual trends or patterns or the exceeding of a threshold value, interpreting results, discussing possible public health action)

- Response
  - Control/response: case management, contact tracing, infection control measures, immunisation activities, improvement of preventive and control measures (vector control, environmental control), community information and education, alerting nearby areas and districts
  - Outbreak investigation: case finding (records, active surveillance), collection and transport of specimens, confirmatory testing, interpretation of results (epidemiological and laboratory)
  - Programme adjustment
  - Changes in policy and planning
- Feedback
- Evaluation and monitoring.

These activities are made possible by a number of support functions that lead to better performance of the core surveillance activities and these should also be assessed:

- Setting standards (e.g., case definitions, standard case management guidelines, standard procedures for investigation)
- Training (surveillance, epidemiology, laboratory)
- Supervision
- Communications systems (e.g. radio, fax, e-mail, phone, health updates)
- Providing resources (human – appropriate number with adequate skills and competencies; material - vehicles, laboratory equipment, supplies etc; financial).

#### **4. Assessing output**

The assessment will provide information on the effectiveness and efficiency of the system(s) in monitoring communicable diseases for prevention and control. The system attributes should be considered (simplicity, flexibility, completeness, sensitivity, timeliness, representativeness). The output of the system (ex. reports) should be able to reflect whether or not the system is achieving its objectives.

#### **5. Integration/Co-ordination/Synergy**

Integration refers to the co-ordination of all surveillance activities and of the support functions common to all control programmes (e.g., data collection, training, and supervision) while leaving follow-up actions to the

different specific intervention programmes. Many functions in the surveillance of most communicable disease are similar and as such offer opportunities for integration. The level of integration/synergy in the national surveillance system can affect the performance, cost and sustainability of the system. Opportunities for integration, synergy and co-ordination should be identified during the assessment for diseases under surveillance.

#### ***6. Laboratories***

Laboratories are essential to disease surveillance and most epidemiological surveillance systems require a laboratory component for confirmation. These serve both for the routine confirmation of clinical syndromes and for rapid confirmation of the causative agent in outbreaks. In some cases the surveillance is completely laboratory-based (example: surveillance of anti-microbial resistance). Assessment of the laboratory capacity (availability, functionality and level of sophistication) should be undertaken in order to determine the role of the laboratory at a given level for surveillance.

#### ***7. Health mapping: the geographic information system (GIS)***

GIS provides an excellent means of collecting and managing epidemiological surveillance and programmatic information. These data can easily be visualised and analysed in a map, showing trends and inter-relationships that would be more difficult to discover in tabular format. GIS allows decision-makers and planners to visualise the health situation of populations easily in relation to the surrounding environment and the existing health and social infrastructures such as health facilities, schools and water supply. Specific diseases and health events can be mapped in relation to the number and location of health facilities, in order to create a comprehensive picture of the health situation of a given community, district or nation. When mapped together, this information creates a powerful tool not only for monitoring surveillance results but also for operational planning and for the targeting interventions and resources to areas/communities in need. This database serves as a common geographic platform within which all surveillance and programmatic data can be concentrated at the most appropriate level. As such GIS constitutes itself as an entry point for integrating disease-specific surveillance approaches.

#### ***8. Communication***

Good communication systems are critical for effective surveillance. In some countries, communication offices are available at varying level of the health care system, with strategic plans, emergency media response plans and trained staff. Others have resources such as computers, appropriate software, with email connections. Many countries use computerized systems for data

collection, reporting, analyzing, feedback and dissemination. Data reported through appropriate electronic system would facilitate the integration of surveillance activities especially if the system is user-friendly, does not use multiple and different data sets that results in extra work load and subsequent abandoning. Radio calls are used in other remote areas. Communication systems should be assessed, taking into account local realities. A description of the communication practices, as well as resources should be made, and needs identified. The outputs of these systems should be assessed (health bulletins, reports, scientific publications, audio/video productions) and the content should be considered (health topics, surveillance data, outbreak investigation, recommendations, etc).

### **What should guide the assessment?**

The procedure proposed in this guideline aims to involve the MoH as the key player in the assessment. The role of the external team is to facilitate the process using standard methods and tools, as recommended by WHO. The end result should be a national plan designed by nationals. This may not result in the perfect plan by external standards but will have a higher chance of success. The goal is to agree on a plan of action (PoA) and to establish a follow-up programme.

The government should accept that, in the long run, surveillance is a core public health function and as such should be funded within the health budget. Political commitment and financial support by the government is essential to obtain sustainable change within the surveillance system if this is to lead towards improvements in disease control. It is important that the solutions to problems are decided by the nationals, and perceived as relevant to the realities within the national health service. External funds from WHO or other donors should be used as a means to get things started in crucial domains.

The procedure should be to involve representatives of the MoH, the individual surveillance focal points for each health event and workers from each level of the system in a facilitated national process.

### **Procedures, activities and timetable of the assessment**

The guideline below outlines a 17 working day (3 weeks) schedule to complete the assessment. This is only a guide since many factors such as the size of the country, the logistics for fieldwork and the availability of senior MoH staff may influence the schedule.

### **Schedule for national surveillance assessment**

|  |                                     |  |
|--|-------------------------------------|--|
| <b>PHASE I*</b><br>Planning                    | Before<br>assessment                | Planning the mission   |
| <b>PHASE II</b><br>Step 1<br>Pre-assessment    | DAYS 1-3                            | Pre-assessment facilitated workshop to examine surveillance priorities and objectives. Further sensitise on the multi-disease approach to surveillance, agree on the list of national priority diseases, adapt the assessment protocol, plan fieldwork |
| Step 2<br>Training                             | DAYS 4-6                            | Training of assessment team members and data managers. Pre-test and adapt assessment tools; finalise logistical requirements, travel to assessment sites   |
| Step 3<br>Field assessment                     | DAYS 7-12                           | Field assessment and travel  |
| Step 4<br>Analysis and report                  | DAYS 13-16                          | Write a preliminary report using a standard format on the assessment findings  |
| Step 5<br>Findings and<br>follow-up schedule   | DAYS 17                             | Post-assessment workshop to present preliminary findings; discuss follow-up schedule and agree to it   |
| <b>PHASE III</b><br>National Plan of<br>Action | After<br>assessment:<br>4 – 8 Weeks | Workshop to elaborate National Plan of Action and implementation framework   |
| <b>PHASE IV</b><br>Follow up                   |                                     | Follow-up implementation of the Plan of Action   |

\*The duration of each phase and step may vary depending on the size of the country.

#### **PHASE I: Planning the mission**

Planning the assessment is essential for the success of the mission. The process begins when a country requests assistance from WHO to carry out an assessment of its communicable disease surveillance system(s). The country is asked to set up a co-ordinating body with a focal person in the MoH and a proposed time frame for the assessment. Key partners including someone from the WHO/WRO should be part of the co-ordinating body. The WHO Country Office should also decide on a counterpart to the MoH focal person.

The WR Office and the MoH should begin work on logistic requirements (transport, lodging, finances, personnel, office facilities and supplies etc) for the assessment (See Annex 2.0. and 2.1. for mission planning spreadsheet and logistic checklist).

Before the assessment a co-ordination meeting should be held between all the external consultants, preferably within the country, together with the WR. This will provide the opportunity of gaining a common understanding of the assessment as well as getting a briefing from the WR about the country. A tentative work plan of the assessment should be drafted, outlining the roles and responsibilities of team members.

It is also crucial to learn about the health and economic system in the country (Recommended documents for reading include WHO, UNAIDS and UNDP Country Profiles as well as Demographic and Health Surveys).

A meeting should be held as soon as possible with the national team. The participation of senior decision-makers at the MoH in all steps of the assessment is critical: if decision-makers are not part of the assessment, the recommendations will not gain the necessary political support within the government. The WHO country representative should therefore ensure this involvement. The WR should assign a focal point in the WHO office to act as liaison before the mission, to take an active part in the process and to follow up on an ongoing basis with the MoH after the assessment. In some countries, the WHO office now has a country epidemiologist who liaises directly with the MoH. It may be useful to have a joint planning sheet for the MOH and WRO (See Annex 2.2).

***Composition of the assessment team: External team  
(Members not resident in the Country)***

The external team should ideally include an epidemiologist, a laboratory expert, a GIS expert, and the designated WHO Country Office focal person. This team may be drawn from the WHO Country Office, the WHO/Regional Office, WHO/HQ and other partners. A team leader should assume overall responsibility for the mission as well as for implementation and follow-up. The external team will facilitate the assessment process and participate in the field assessment. In collaboration with the national team leader, the external team leader will coordinate the assessment process, including the writing of the assessment report. Everyone should be familiar with the Terms of Reference (TOR) for the assessment (See Annex 3.1 for prototype TOR).

***National team***

The national team shall be drawn from various levels of the health services and from all major disease control programmes, national institutions such as Field Epidemiology Training Programmes (FETPs) and NGOs. Broad national representation will ensure a more equitable assessment and allow the various players to interact professionally. It is essential that all team members be briefed on the objectives of the assessment. The MoH shall designate a national counterpart to the external team leader and a focal person who will liaise with the WHO focal person.

## PHASE II: The assessment

### Step 1: Pre-assessment facilitated workshop with national team

A courtesy visit to the Minister of Health should take place, to brief her/him on the objectives of assessment before the workshop takes place.

The aim of the workshop is to take the group through a process of examining disease priorities and surveillance objectives, agreeing on the protocol and adapting generic tools for the field assessment of surveillance system(s) performance. The workshop includes several activities, each of which leads to a product that may be used for the next activity. The activities themselves are part of assessment and the product of each session will provide useful information for the final report. The workshop usually lasts 3 days. The starting and finishing times for each day should be determined by the local working day.

#### **Activities and products from pre-assessment workshop**

| Activity  | Products   |
|---|--|
| 1. Plenary session on the multi-disease approach and the objectives of assessment (Annexes 3)             | 1.1 MoH decision-makers sensitised on the multi-disease approach and on assessment objectives  |
| 2. Exercise: setting priorities for communicable diseases (Annex 4)                                       | 2.1 Adoption of list of Priority communicable diseases   |
| 3. Inventory of current surveillance activities (Annex 5)   | 3.1 Table summarizing all current surveillance activities  |
| 4. Surveillance objectives and indicators (Annex 6)   | 4.1 A table summarizing surveillance objectives and indicators for each priority disease under surveillance  |
| 5. Surveillance process and task description, by health service level (Annex 7)                           | 5.1 Flow diagrams to illustrate surveillance process<br>5.2 Table for each priority disease showing the tasks that are carried out at each level of the system |
| 6. Adaptation of tools for field assessment (Annex 8 and 13)  | 6.1 Indicators to test system performance<br>6.2 Checklist/questionnaires for data collection  |
| 7. Selection of assessment sites, finalisation of teams, organization, and scheduling of visits (Annex 9) | 7.1 Sample sizes and map showing districts and facilities to be visited<br>7.2 Table showing organization of each team, sites to be visited, and timing        |
| 8. Logistics for field visits (Annex 9.1)   | 8.1 Table showing transport, security, accommodation, financial and administrative arrangements for the team   |

## Step 2: Training of assessment teams

Training of the assessment team is a continuation of the facilitated workshop and comprises consensus building, pre-testing and revision of the tool. During this training session, team members are expected to examine the data collection tools and get a clear and common understanding of the questions and of what exactly to look for while conducting the interview. The training should include a demonstration of various sample analyses. The team leader moderates the training sessions in collaboration with the national counter part.

The content of the training is as follows:

- Conduct during field visit
- Information meeting with local team
- Detailed organization of assessment
- Data collection process: questionnaire use (quality control)
- Data entry, cleaning of data and draft analysis
- Field testing, feedback and adaptation of the assessment tools.

### **Activities and products from training workshop**

| <b>Activities</b>   | <b>Products</b>  |
|---|--|
| Briefing on expectations on arrival and contacts with local authorities on site   | Conduct (see Annex 9.3) and administrative arrangements known                        |
| Information meeting with local team   | Content and conduct of the meeting mastered  |
| Detailed organization of the assessment (Role of team members, number and types of sites for assessment, tracking questionnaires, identification of interviewees, appointments, transport, security, accommodation etc) | Detailed organization of assessment known  |
| Data collection process: checklist/questionnaire use (filling, quality control)   | 1. Questions understood<br>2. Data collection mastered                               |
| Data entry, cleaning and draft analysis   | 1. Capacity built for data entry and cleaning<br>2. Draft analysis programme adopted |
| Field testing, feedback and adaptation of the assessment tools  | 1. Assessment tools field-tested<br>2. Assessment tools adapted                      |

## Step 3: Field assessment

The main aim of the field visits is to gather information on the pre-designed tools to carry out a formal assessment of the performance for all components of the surveillance system. The field assessment should last 3 to 7 days.

Advance arrangements and planning are critical to the success of this step. Preparations for the field visits should be made by the MoH with the support of the WHO office, prior to the arrival of the assessment team.

The site visits should be carried out according to an agreed timetable; they may involve a team visiting both peripheral and intermediate levels. Each type of site visited will require a specific checklist/questionnaire. Working with the tools developed will involve asking questions, observing practices and gathering documentation of activity.

The approach at each site visited shall be to:

- Have an initial meeting to introduce the objectives of the assessment and to ask relevant questions
- Obtain informal feedback on problems and issues that workers themselves have identified regarding surveillance
- Identify examples of good and bad practice
- Consult reports of outbreaks or other investigations
- Make sure that checklists/questionnaires are filled in legibly
- Record and if possible resolve any problems or ambiguities in the tools
- Clean data
- Enter data into a pre-prepared database.

The assessing team should meet regularly at the end of the day or once every two days to document the problems encountered, the challenges, strengths and weakness of the sites visited, the systems assessed, the laboratory linkages to surveillance etc. This qualitative analysis would contribute to the interpretation of the quantitative analysis.

#### **Step 4: Analysis and preliminary report writing**

Writing the report should be a team activity, usually lasting 3 days and involving:

- Analysis of the products of the pre-assessment workshop
- Analysis of data from the field visits, both qualitative (impressions obtained during the visits) and quantitative (replies to questionnaires)
- Identification of strengths, weaknesses, opportunities and threats in the national surveillance and response system
- Identification of solutions, opportunities, threats to integration

- Recommendations to strengthen the capacity, improve co-ordination, build synergies, and take advantage of driving forces for the national surveillance and response system.

The assessment report (see prototype in Annex 11.1) should use the standard surveillance terms provided in Annex 1.

The report should refer to the priority diseases and to capacity and co-ordination/integration of the surveillance system(s).

#### ***Priority Diseases***

Are current surveillance activities adequate in terms of the diseases covered and the population under surveillance? The revised list of priority diseases should be included.

#### ***Capacity***

For this section the capacity should refer to the performance of the core surveillance activities and the surveillance support functions. Field visits will be the source of this information and as such this section will reflect the surveillance methods.

Analysis of capacity may be undertaken for:

- All diseases
- Indicator diseases (e.g. measles for EPI, gonorrhoea for STIs and cholera for epidemic-prone diseases)
- Groups of diseases (e.g. vaccine-preventable diseases in EPI).

This will depend on how many diseases were included in the field assessment.

#### ***Co-ordination/Integration***

The level of co-ordination/integration should be reported in terms of the core functions and support functions. Do disease surveillance systems/control programmes use the same mechanisms to carry out any of the functions and what are the areas where further synergy would be beneficial?

### **Step 5: Post-assessment workshop to present preliminary findings**

A major challenge in strengthening surveillance systems is the actual implementation of change. One of the most difficult tasks in surveillance assessment and strengthening is to transform a report with an implementation plan into real activities over a period of time. One way of doing this is:

- To get political commitment into the process
- To get the MoH to commit resources to the process
- To identify critical activities that would benefit from outside technical support
- To follow up on all commitments systematically and ensure a co-ordinated implementation process.

To this end, a one-day workshop at the end of the assessment may prove invaluable in bringing together decision-makers from the different parties and stakeholders in order to obtain a clear agreement on the activities to be carried out and supported. These activities should have a timetable and identify responsible individuals and resources.

Attendance at the end of assessment workshop should include:

- Ministry of Health
- World Health Organization
- Donors (e.g., ADB, USAID, EU, DANIDA, DFID)
- Other UN agencies (e.g., UNDP, UNICEF)
- Others partners (e.g., CDC, NGOs, academic institutions, representatives of private practitioners)
- Laboratory Institutions outside the Ministry of Health.

The workshop should include the following way:

1. Presentation of the draft report by the assessment team
2. Discussion of the assessment findings
3. Agreement on future activities (i.e., timeline for the final assessment report and Plan of Action workshop)
4. Consensus of all stakeholders to consider the implications of the assessment findings and recommendations in the execution of their duties and in their surveillance strengthening efforts.

### **PHASE III: Workshop to elaborate plan of action**

The workshop should take place 1-2 months after the assessment. During this time, the preliminary report should be finalised and circulated to all concerned.

Participants coming from all levels of the health system (central, intermediate and district including health facilities) should elaborate a draft plan of action. This working group should:

- Prepare a draft implementation plan and agree on activities and budget
- Agree on the final implementation plan with a prioritised list of activities and proposed timetable and an allocation of responsibilities
- Agree on follow-up method and schedule.

The implementation plan should centre on priority activities that can improve the surveillance and response systems (see PoA matrix Annex 13). This plan will be presented at a one-day session on the last day of the workshop for discussion and approval.

The implementation plan should:

- Identify priority activities
- Set timetables for the activities
- Identify the person or agency responsible for each activity and for overall implementation
- Estimate costs
- Identify what percentage of the costs are to be borne by the Government
- Identify indicators of activity implementation and success
- Suggest a process of formal follow-up and evaluation of implementation both
  - Routinely through an update/monitoring tool
  - Formally through a follow-up evaluation at least once a year.

Attendance at the final session of the Plan of Action workshop should include:

- Ministry of Health
- World Health Organization

- Donors (e.g. ADB, USAID, EU, DANIDA, DFID)
- Other UN agencies (e.g. UNDP, UNICEF)
- Other partners (e.g. CDC, NGOs, academic institutions, representatives of private practitioners)
- Laboratory Institutions outside the Ministry of Health.

#### **PHASE IV: Follow-up of the implementation of the multi-disease approach to surveillance**

Follow-up is critical to the success of the process. The MoH should provide regular standardised updates on the progress and on the problems encountered. WRO will send regular progress reports to the Regional Office/Head Quarters.

WHO and partners will carry out an external evaluation of the implementation of the surveillance and response strengthening efforts, as well as the multi-disease approach. It is suggested that a midterm (2<sup>nd</sup> to 3<sup>rd</sup> year) review and a 5-year external review of the progress of implementation of the objectives in the Action Plan should be undertaken. Internal (in-country) reviews should be undertaken annually.

# LIST OF ANNEXES

---

|            |   |     |
|------------|---|-----|
| ANNEX 1.0  | SURVEILLANCE DEFINITIONS .....  | 21  |
| ANNEX 2.0  | MISSION PLANNING CHECKLIST .....  | 29  |
| ANNEX 2.1  | LOGISTICS CHECKLIST .....   | 31  |
| ANNEX 2.2  | MOH/WR PLANNING SPREADSHEET .....   | 33  |
| ANNEX 3.0  | SETTING OF OBJECTIVES FOR ASSESSMENT AND TEAM ORGANIZATION.....             | 35  |
| ANNEX 3.1  | PROTOTYPE TERMS OF REFERENCE FOR ASSESSMENT .....                           | 37  |
| ANNEX 3.2  | LIST OF PARTICIPANTS IN ASSESSMENT TEAM.....                                | 39  |
| ANNEX 4.0  | PRIORITY SETTING EXERCISE .....   | 41  |
| ANNEX 5.0  | INVENTORY OF CURRENT SURVEILLANCE ACTIVITIES .....                          | 43  |
| ANNEX 5.1  | DESCRIPTION OF MAJOR SURVEILLANCE ACTIVITIES<br>TO IDENTIFY GAPS .....      | 45  |
| ANNEX 6.0  | SURVEILLANCE SYSTEM(S), FLOW CHART(S) AND TASK DESCRIPTION .....            | 47  |
| ANNEX 7.0  | TASK ANALYSIS BY LEVEL FOR PRIORITY DISEASES.....                           | 49  |
| ANNEX 8.0  | DESIGNING TOOLS FOR FIELD ASSESSMENT .....                                  | 51  |
| ANNEX 9.0  | SELECTION OF ASSESSMENT SITES AND SCHEDULING OF VISITS.....                 | 53  |
| ANNEX 9.1  | SELECTION OF SAMPLES FOR REGIONS, DISTRICTS,<br>AND HEALTH FACILITIES.....  | 55  |
| ANNEX 9.2  | SCHEDULE AND LOGISTICS.....   | 57  |
| ANNEX 9.3  | CONDUCT DURING FIELD ASSESSMENT.....  | 59  |
| ANNEX 10.0 | ANALYSIS, PRELIMINARY REPORT WRITING .....                                  | 61  |
| ANNEX 10.1 | PROTOTYPE REPORT WRITING FORMAT .....                                       | 63  |
| ANNEX 11.0 | POA MATRIX .....  | 67  |
| ANNEX 12.0 | GENERIC QUESTIONNAIRES.....   | 69  |
| ANNEX 13.0 | LABORATORY ASSESSMENT .....   | 95  |
| ANNEX 14.0 | ASSESSMENT OF GEOGRAPHIC INFORMATION SYSTEMS<br>AND MAPPING RESOURCES ..... | 111 |



## SURVEILLANCE DEFINITIONS

---

These definitions are standardized by WHO and as such are referred to in the guidance below. All reports to and by WHO should preferably use these terms as defined in this glossary in order to improve standardization.

**ACCEPTABILITY** Acceptability is measured by the willingness of persons conducting surveillance and those providing data to generate accurate, consistent and timely data.

**ACTIVE CASE FINDING** The process of seeking out cases or health event under surveillance (e.g. house visits by community workers to identify cases of tuberculosis, active searching of medical records to identify cases of acute haemorrhagic fever).

**ATTACK RATE** The cumulative incidence of infection in a group observed over a period during an epidemic. This “rate” can be determined empirically by identifying clinical cases and/or by means of seroepidemiology. Because its time dimension is uncertain or arbitrarily decided, it should probably not be described as a rate. (*Last JM, A Dictionary of Epidemiology, 2001*).

**CARRIER** A person or animal that harbours a specific infectious agent in the absence of discernible clinical disease and serves as a potential source of infection. The carrier state may occur in an individual with an infection that is inapparent throughout its course (known as healthy or asymptomatic carrier) or during incubation period, convalescence, and post convalescence of an individual with a clinically recognisable disease (known as incubatory carrier or convalescent carrier). The carrier state may be of short or long duration (temporary or transient carrier or chronic carrier). (*Last JM, A Dictionary of Epidemiology, 2001*).

**CASE** A person who has the particular disease, health disorder, or condition which meets the case definitions for surveillance and outbreak investigation purposes. The definition of a case for surveillance and outbreak investigation purpose is not necessarily the same as the ordinary clinical definition. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**CASE CLASSIFICATION** Gradations in the likelihood of being a case (e.g., suspected / probable / confirmed). This is particularly useful where early reporting of cases is important (e.g., Ebola haemorrhagic fever) and where there are difficulties in making definite diagnoses (e.g., specialized laboratory tests required).

**CASE DEFINITION** A set of diagnostic criteria that must be fulfilled for an individual to be regarded as a case of a particular disease for surveillance and outbreak investigation purposes. Case definitions can be based on clinical criteria, laboratory criteria or a combination of the two with the elements of time, place and person.

**CASE-FATALITY RATE** The proportion of cases of a specified condition which are fatal within a specified time. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

$$\text{Case-fatality rate} = \frac{\text{Deaths from a given disease in a given period} \times 100}{\text{Diagnosed cases of that disease (in the same period)}}$$

**CLUSTER** Aggregation of relatively uncommon events or diseases in space and/or time in amounts that are believed or perceived to be greater than could be expected by chance. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**COMMUNICABLE DISEASE (SYNONYM: INFECTIOUS DISEASE)** An illness due to a specific infectious agent or its toxic products that arises through transmission of that agent or its products from an infected person, animal, or reservoir to a susceptible host, either directly or indirectly through an intermediate plant or animal host, vector, or the inanimate environment. (*Last JM, ed. A Dictionary of Epidemiology, 2001*).

**CONTACT (OF AN INFECTION)** A person or animal that has been in such association with an infected person or animal or a contaminated environment as to have had opportunity to acquire the infection. (*Last JM, A Dictionary of Epidemiology, 2001*).

**CONTACT TRACING** see active case finding.

**EARLY WARNING SYSTEM** In disease surveillance, a specific procedure to detect as early as possible any abnormal occurrence or any departure from usual or normally observed frequency of phenomena (e.g. one case of Ebola fever). An Early Warning System is only useful if linked to mechanisms for early response. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**ELIMINATION** Reduction of case transmission to a predetermined very low level; e.g., elimination of tuberculosis as a public health problem was defined by the WHO (1991) as reduction of prevalence to a level below one case per million population. (*Last JM, A Dictionary of Epidemiology, 2001*).

**EMERGING INFECTIONS** A collective name for infectious diseases that have been identified and taxonomically classified recently. In the final quarter of the twentieth century, more than 30 such conditions, many of them capable of causing dangerous epidemics, were recognized. They include human immuno-deficiency virus (HIV) infection, ebola virus disease, hantavirus pulmonary syndrome and other viral haemorrhagic fevers, campylobacter infection, transmissible spongiform encephalopathies, legionnaires' disease, and lyme disease. Some appear to be "new" diseases of humans, others may have existed for many centuries and have been recognized only recently because ecological or other environmental changes have increased the risk of human infection. re-emerging infections are certain "old" diseases, such as tuberculosis and syphilis, that have experienced a resurgence because of changed host-agent-environment conditions. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**ENDEMIC** The constant presence of a disease or infectious agent within a given geographic area or population group; may also refer to the usual prevalence of a given disease within such area or group. The expression "endemic disease" has a similar meaning. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**EPIDEMIC** [from the Greek επι (upon), δῆμος (people)]. The occurrence in a community or region of cases of an illness, specific health-related behaviour, or other health-related events clearly in excess of normal expectancy. The community or region and the period in which the cases occur are specified precisely. The number of cases indicating the presence of an epidemic varies according to the agent, size, and type of population exposed, previous experience or lack of exposure to the disease, and time and place of occurrence. (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**EPIDEMIC THRESHOLD** The number or density of susceptibles required for an epidemic to occur. (e.g. meningococcal meningitis: see exception flagging system). (*Adapted from Last JM, A Dictionary of Epidemiology, 2001*).

**EXCEPTION FLAGGING (REPORTING) SYSTEM** A manual or automated system of data analysis which calculates thresholds for epidemic or outbreak detection (e.g. the signal given when incidence of meningococcal meningitis in African belt area is 15/100 000/week over 2 consecutive weeks).

**EXPOSURE** Proximity and/or contact with a source of a disease agent in such a manner that effective transmission of the agent, harmful or protective effects of the agent may occur. (*Adapted from Last JM, ed. A Dictionary of Epidemiology, 2001*).

**FEEDBACK** The regular process of sending analyses and reports about the surveillance data back through all levels of the surveillance system so that all participants can be informed of trends and performance.

**FLEXIBILITY** Flexibility is a measure of the ability of the surveillance system to be easily adapted to new reporting needs in response to changes in the nature or the importance of the health event, the population monitored, or the resources available.

**GENERALIZABILITY/VALIDITY/REPRESENTATIVENESS** The degree to which inference can be drawn from the information gathered by the surveillance system to the target population.

**GIS** An organized collection of computer hardware, software, geographical data and personnel designed to efficiently capture, store, update, manipulate, analyse and display all forms of geographically referenced information. It is first and foremost an information system with a geographical variable, which enable users to easily process, visualize and analyse data or information spatially. GIS can be used to prepare models showing trends in time and space. Satellite imaging and remote sensing have expanded its scope (e.g. to identify regions prone to malaria).

**HEALTH EVENT** Any event relating to the health of an individual (e.g., the occurrence of a case of a specific disease or syndrome, the administration of a vaccine or an admission to hospital).

**INCIDENCE** The number of instances of illness commencing, or of persons falling ill, during a given period in a specified population. (*Prevalence and Incidence. WHO Bulletin, 1966, 35: 783-784*).

**INCIDENCE RATE** The rate at which new events occur in a population. The numerator is the number of new events that occur in a defined period; the denominator is the population at risk of experiencing the event during this period, sometimes expressed as person-time. (*Adapted from Last JM, ed. A Dictionary of Epidemiology, 2001*).

**INFECTIOUS DISEASE SEE COMMUNICABLE DISEASE**

**NOTIFIABLE DISEASE** A disease that, by statutory/legal requirements, must be reported to the public health or other authority in the pertinent jurisdiction when the diagnosis is made. (*Adapted from Last JM, ed. A Dictionary of Epidemiology, 2000*).

**NOTIFICATION** The processes by which cases or outbreaks are brought to the knowledge of the health authorities. In the context of the *International Health Regulations*, notification is the official communication of a disease/health event to the World Health Organization by the health administration of the Member State affected by the disease/health event.

**OUTBREAK** An epidemic limited to localised increase in the incidence of a disease, e.g., in a village, town, or closed institution. (*Adapted from Last JM, ed. A Dictionary of Epidemiology, 2001*).

**PERFORMANCE INDICATORS** Specific agreed measurements of how participants are functioning within the surveillance or reporting system. These indicators may measure both the process of reporting (e.g., completeness, timeliness ) and the action taken in response to surveillance information (e.g., the percentage of cases investigated or surveyed) and the impact of surveillance and control measures on the disease or syndrome in question (e.g., the percentage of outbreaks detected by the system, the drop in the number of cases over a specified time period).

**PERIODICITY** A repeating pattern of a phenomenon or an event, especially the repetition of comparable values, e.g., seasonal fluctuation in numbers of cases of respiratory infections. (*Last JM, A Dictionary of Epidemiology, 2001*).

**PREVALENCE** The number of instances of illness or of persons ill, or of any other event such as accidents, in a specified population, without any distinction between new and old cases. Prevalence may be recorded at a stated moment (point prevalence) or during a given period of time (period prevalence). (*Prevalence and Incidence. WHO Bulletin, 1966; 35:783-784*).

**PREVALENCE RATE** The total number of all individuals who have an attribute or disease at a particular time (or during a particular period) divided by the population at risk of having the attribute or disease at this point in time or midway through the period. (*Last JM, A Dictionary of Epidemiology, 2001*).

**REPORTING COMPLETENESS** Proportion of all expected reports that were actually received. It is usually stated as “% completeness as of a certain date” (e.g. if of 30 administrative units in a reporting system 15 submit reports, the reporting completeness is 50%; if of 50 cases of diarrhoea 40 are reported, the reporting completeness is 80%).

**REPORTING SYSTEM** The specific process by which diseases or health events are reported. This will depend on the importance of the disease and the type of surveillance.

**REPORTING TIMELINESS** Proportion of all expected reports in a reporting system received by a given date (due date).

**SECULAR TREND** (Synonym: temporal trend) Changes over a long period of time, generally years or decades. (*Adapted from Last JM, ed. A Dictionary of Epidemiology, 2001*).

**SEROSURVEILLANCE** The surveillance of an infectious disease through immunological markers of the disease in a population or sub-population (e.g. measuring the presence of HIV antibodies in pregnant women coming for antenatal care).

**SENSITIVITY IN SURVEILLANCE** The ability of a surveillance or reporting system to detect true health events i.e. the ratio of the total number of health events detected by the system over the total number of true health events as determined by an independent and more complete means of ascertainment.

**SPECIFICITY IN SURVEILLANCE** A measure of how infrequently a system detects false positive health events i.e. the number of individuals identified by the system as not being diseased or not having a risk factor, divided by the total number of all persons who do not have the disease or risk factor of interest.

**SURVEILLANCE** The process of systematic collection, orderly consolidation and evaluation of pertinent data with prompt dissemination of the results to those who need to know, particularly those who are in a position to take action (*Adapted from Report of the Technical Discussions at the twenty-first World Health Assembly on National and Global Surveillance of Communicable Diseases, 18 May 1968 — A21/Technical Discussion/5*).

**SURVEILLANCE, ACTIVE** Surveillance where public health officers seek reports from participants in the surveillance system on a regular basis, rather than waiting for the reports (e.g. telephoning each participant monthly).

**SURVEILLANCE, CASE-BASED** Surveillance of a disease by collecting specific data on each case (e.g. collecting details on each case of acute flaccid paralysis (AFP) in poliomyelitis surveillance).

**SURVEILLANCE, COMMUNITY** Surveillance where the starting point for the notification is from community level, normally reported by a community worker. It can be active (looking for cases) or passive (reporting cases). This may be particularly useful during an outbreak and where syndromic case definitions can be used (the active identification of community cases of Ebola virus infection in Kikwit was an example of active community surveillance).

**SURVEILLANCE, ENHANCED** The collection of additional data about cases reported under routine surveillance. Routine surveillance is a starting point for more specific data collection on a given health event. This information may be sought from the reporter, the case, and the laboratory or from another surveillance data set.

**SURVEILLANCE, HOSPITAL-BASED** (Synonym: Hospital surveillance) Surveillance where the starting point for notification is the identification by a hospital of a patient with a particular disease or syndrome.

**SURVEILLANCE, INTENSIFIED** The upgrading from a passive to an active surveillance system for a specified reason and for a limited period (usually because of an outbreak ). It must be noted that the system then becomes more sensitive; secular trends may therefore need to be interpreted carefully.

**SURVEILLANCE, LABORATORY** Surveillance where the starting point is the identification or isolation of a particular organism in a laboratory (e.g. surveillance of salmonellosis).

**SURVEILLANCE, PASSIVE** Surveillance where reports are awaited and no attempts are made to seek reports actively from the participants in the system.

**SURVEILLANCE, ROUTINE** The regular systematic collection of specified data in order to monitor a disease or health event.

**SURVEILLANCE, SENTINEL** Sentinel surveillance is surveillance based on the collection of data from a sample (random or non-random) of collecting sites as indicator data for the rest of the population, in order to identify cases of a disease early or to obtain indicative data about trends of a disease or health event. Examples are the use of a few hospitals to monitor the composition of influenza virus and check that the vaccine includes the right components, or the use of a network of general practitioners to monitor diseases or health events (e.g. attempted suicide, requests for HIV testing). One instance of sentinel surveillance is the use of a particular population group (e.g., monitoring the serology of syphilis or HIV infection among pregnant women as an indicator of trends in the general population). Sentinel surveillance is inappropriate for those situations where every case requires public health action, e.g., poliomyelitis.

In sentinel surveillance standard case definitions and protocols must be used to ensure validity of comparisons across time and sites despite lack of statistically valid sampling. Sentinel surveillance may include the use of animal sentinels to detect circulation of arboviruses.

**SURVEILLANCE REPORT** A regular publication with specific information on the disease under surveillance. It should contain updates of standard tables and graphs as well as information on outbreaks etc. In addition it may contain information on the performance of participants using agreed performance indicators.

**SURVEY** An investigation in which information is systematically collected. Usually carried out in a sample of a defined population group, within a defined time period. Unlike surveillance it is not ongoing; however, if repeated regularly, surveys can form the basis of a surveillance system.

**SYNDROME** A symptom complex in which the symptoms and/or signs coexist more frequently than would be expected by chance on the assumption of independence. (*Last JM, ed. A Dictionary of Epidemiology, 2001*).

**SYNDROMIC REPORT** The notification of a health event under surveillance for which the case definition is based on a syndrome not on a specified disease (e.g. acute haemorrhagic fever syndrome, acute respiratory syndrome).

**ZERO REPORTING** The reporting of “zero case” when no cases have been detected by the reporting unit. This allows the next level of the reporting system to be sure that the participant has not sent data that have been lost, or that the participant has not forgotten to report.

## MISSION PLANNING CHECKLIST

| <b>Task</b>  | <b>Responsibility</b>                     |
|--|---|
| Ensure MoH commitment to the process   | WHO Country Office                        |
| Get a clear briefing from WHO to all members of the team from WHO on the objectives of the mission               | WHO Regional Office<br>WHO HQ             |
| Make sure the team leader is clearly identified  | WHO Regional Office                       |
| Make sure the WR for the country is fully informed and involved  | WHO Regional Office                       |
| Make sure the necessary invitations are sent   | WHO Regional Office                       |
| Identify a focal point person within the Country Office  | WHO Country Office                        |
| Find out about the country, the health services and the surveillance system(s)                                   | WHO Country Office<br>WHO Regional Office |
| Send background WHO assessment material to WRO and MoH   | WHO HQ<br>WHO Regional Office             |
| Ensure that the MoH is well briefed / sensitised by the WR on the multi-disease approach to disease surveillance | WHO Country Office                        |
| Specify the profile that the assessment participants should fulfil   | WHO Regional Office<br>WHO HQ             |
| Ensure senior representation on the national team  | WHO Country Office<br>MoH                 |
| Ensure representation from various levels of the system and from all major control programmes                    | WHO Country Office<br>MoH                 |
| Identify a focal point person in the MoH for planning and carrying out the assessment                            | MoH                                       |
| Prepare logistic arrangements for the mission  | WHO Country Office<br>MoH                 |
| Identify a venue for the workshop  | WHO Country Office<br>MoH                 |
| Organize access to computers, printers, photocopiers and secretarial services                                    | WHO Country Office<br>MoH                 |
| Arrange travel and accommodation arrangements as appropriate   | WHO Country Office                        |



## LOGISTICS CHECKLIST

---

### **Arrival in country**

- WR arranges reception at airport, visa provisions (if required)
- WR arranges hotel reservation
- WR arranges security clearance if necessary.

### **Personnel**

- WR/MOH designates administrative and secretarial staff
- MoH makes administrative arrangements for participation of national staff.

### **Office facilities**

- WR/MOH arranges office facilities including communication for the assessment team
- WR and MoH arrange for the workshop site and equipment.

### **Transport**

Transport arranged by MoH and WR.

### **Other**



ANNEX 2.2

# MOH/WR PLANNING SPREADSHEET

**Spreadsheet for the planning of surveillance assessment**

| Country   | Dates of assessment             |                           |                  |          |
|---|---------------------------------|---------------------------|------------------|----------|
| Task  | Person Responsible<br>Name/Unit | Expected date<br>DD/MM/YY | Completed<br>Y/N | Comments |
| Discuss mission with MoH  |                                 |                           |                  |          |
| Obtain country clearance and invitation   |                                 |                           |                  |          |
| Identify external (WRO) and internal (MoH) focal point                          |                                 |                           |                  |          |
| Obtain background material on country health services, surveillance system etc. |                                 |                           |                  |          |
| Share background assessment material MoH  |                                 |                           |                  |          |
| Start logistic arrangements for the mission                                     |                                 |                           |                  |          |
| Meet with relevant donor and technical partners                                 |                                 |                           |                  |          |



## SETTING OF OBJECTIVES FOR ASSESSMENT AND TEAM ORGANIZATION

---

|                      |   |
|----------------------|---|
| Activity:            | Plenary session on the objectives of assessment and finalization of team organization   |
| Objective:           | To finalize the objectives of the assessment and the Organization of the assessment team  |
| Method:              | Group discussion  |
| Duration:            | 1¾ hours  |
| Materials required:  | Prototype terms of reference and prototype team table for the organization of the team  |
| Role of facilitator: | To ensure that the objectives of assessment are established, taking into account specific aspects of the system such as the system of communication, laboratory, GIS, and others that might require special attention |
| Product:             | Agreed terms of reference and table showing Organization of assessment teams  |

### **Step I**

The participants should agree on the objectives of the assessment, keeping in mind that the final report will relate closely to these terms of reference. The methods to be used should be agreed upon, as well as the anticipated outputs (for example, comprehensive documentation of the surveillance system, action plan etc.). The various institutions taking part in the surveillance assessment should be identified.

### **Step II**

The professional role of each team member from participating institutions should be stated, in order to allocate tasks rationally and fairly. The team will take an active part in all aspects of evaluation, and liaise with the various units and organization involved, including following-up assessment after the mission. (See Annex 3.2)

### **Step III**

The relevant details should be filled, using the templates provided or an adapted version thereof (Annex 3.1: Prototype Terms of Reference for Assessment and Annex 3.2: List of Participants in Assessment Team).

#### ***Work plan***

---

---

| <b>Step</b> | <b>Specific task</b>                           | <b>Person responsible</b>    | <b>Duration</b> | <b>Resources</b>                            | <b>Output</b>   |
|-------------|--|------------------------------|-----------------|---|---|
| Step I      | Define objectives and outputs of evaluation    | Team-members                 | 45 minutes      | Background materials on assessment mission  | Record of objectives and expected outputs             |
| Step II     | Elaborate terms of reference                   | Team-members/<br>facilitator | 30 minutes      | Draft ToR                                   | Record of elaborated Terms of Reference               |
| Step III    | Attribute groups and functions to team members | Team-members/<br>facilitator | 30 minutes      | List of team members and professional roles | Table showing the organization of the assessment team |

---

---

## PROTOTYPE TERMS OF REFERENCE FOR ASSESSMENT

---

The Ministry of Health of [COUNTRY] invites the World Health Organization to facilitate the assessment of the national communicable disease surveillance, epidemic preparedness and response with the following objectives:

- To assess the structure, process, capacity, resources, effectiveness and co-ordination of the national surveillance system for communicable diseases, epidemic preparedness and response; and
- To propose a plan of action to strengthen communicable disease surveillance, epidemic preparedness and response.

The assessment will take the form of a facilitated workshop to examine the current system and adapt the generic tool, followed by training of interviewers and by pre-testing. The field assessment will be conducted in sites selected from all levels of the health system. After the field assessment, all relevant findings will be summarised in a report that will identify the strengths and weaknesses of the current system. This report will be presented at a final workshop at which a draft plan of action will be drawn, including agreement on activities, time-tables and budgets.

The assessment team will be led jointly by [NAME] from the Ministry of Health and [NAME] nominated by the World Health Organization. The team itself will consist of Ministry of Health staff from all major control programmes and from the epidemiology unit in the ministry, and WHO staff.







## PRIORITY SETTING EXERCISE

---

|                      |   |
|----------------------|---|
| Objective:           | To categorise relevant communicable diseases according to their public health priority                            |
| Method:              | Small group discussion (8-10 persons per group)   |
| Duration:            | Approximately 2 hours   |
| Materials required:  | Background information on communicable diseases in the country  |
| Role of facilitator: | To help the group complete a template table through examination of background material and small group discussion |
| Product:             | Table of priority communicable diseases with justification  |

### **Step I**

The facilitator should get the group to make a list of criteria to prioritise diseases (high mortality, high morbidity, high case fatality rate, for elimination or eradication, control is feasible, the cost involved, epidemic potential, existing control programmes, national, regional and global targets, etc.) and a list of diseases that should be under surveillance.

### **Step II**

The facilitator should obtain a list of diseases under surveillance in the country.

### **Step III**

These lists should be compared to achieve consensus on what should be under surveillance. Where there is no consensus, the facilitator should assist in a process of prioritisation.



## INVENTORY OF CURRENT SURVEILLANCE ACTIVITIES

---

Using the consensus list of priority diseases, the group should examine the strategies used in the surveillance of these diseases and identify gaps in surveillance if any.

|                      |   |
|----------------------|---|
| Objective:           | To make an inventory of current surveillance activities for the diseases on the consensus list and identify gaps    |
| Method:              | Small group discussion  |
| Duration:            | Approximately 2 hours   |
| Materials required:  | Consensus list of diseases from previous exercise and information on current surveillance activities in the country |
| Role of facilitator: | To help the group complete a template table by examination of background material                                   |

### **Step I**

The facilitator should assist the group in identifying gaps in the existing surveillance system. For each disease questions should be asked about how surveillance is conducted: (see Annex 5.1)

### **Step II**

Participants should produce a consensual document on the model of Annex 5.1

| Step    | Specific task  | Person responsible       | Duration | Resources                          | Output   |
|---------|--|--------------------------|----------|------------------------------------|--|
| Step I  | Inventory of surveillance activities   | Team-members             | 1 hour   | Background on surveillance systems | List of existing surveillance activities and diseases under surveillance |
| Step II | Identification of gaps in the surveillance of the priority diseases identified | Team-members/facilitator | 1 hour   | Template table                     | Table illustrating gaps in surveillance                                  |



## DESCRIPTION OF MAJOR SURVEILLANCE ACTIVITIES TO IDENTIFY GAPS

| Priority Disease | Existing surveillance activities | Programme(s) managing surveillance activities  | Levels at which surveillance activities are carried out | Types of Surveillance |  |                                  |                               | Case Definition | Confirmation (Clinical/lab or both) | Identified gaps |
|------------------|----------------------------------|--|---|-----------------------|--|----------------------------------|-------------------------------|-----------------|-------------------------------------|-----------------|
|                  |                                  |  |   | Collection strategy   | Type of data collected                 | Frequency of collection          | Method of collection          |                 |                                     |                 |
| Cholera          | Y                                | 1. Epidemiological surveillance unit<br>2. CDD | All (Central, District, Health Facility)                | Routine               | Case-based<br>Aggregated               | Immediately<br>Weekly            | Active<br>Passive             | Y               | Both                                |                 |
| VHF              | N                                | None   | None  | Survey                | Case-based                             | Immediately                      | Active                        | N               | Lab                                 |                 |
| HIV              | Y                                | HIV  | All   | Sentinel              | Case-based                             | Annually                         | Passive                       | Y               | Lab                                 |                 |
| Malaria          | Y                                | Epidemiological surveillance unit              | All   | Routine Surveys       | Aggregated<br>Case-based<br>Case-based | Monthly<br>Annually<br>2-5 years | Passive<br>Passive<br>Passive | Y               | Both                                |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |
|                  |                                  |  |   |                       |  |                                  |                               |                 |                                     |                 |



## SURVEILLANCE SYSTEM(S), FLOW CHART(S) AND TASK DESCRIPTION

Using the consensus list of priority diseases the group should study the design of the surveillance systems and the process by which data and samples move through the system. The group should also identify those units responsible for response and feedback.

|                      |   |
|----------------------|---|
| Objective:           | To draw a flow chart showing design of surveillance system and task description by level                      |
| Method:              | Group discussion  |
| Duration:            | Approximately 3 hours   |
| Materials required:  | Products from previous session and any documentation of current systems                                       |
| Role of facilitator: | To help the group to produce the flow chart by examining the background material and through group discussion |
| Product:             | Annotated flow chart(s)   |

### **Work plan**

| Step    | Specific task  | Person responsible       | Duration | Resources   | Output   |
|---------|--|--------------------------|----------|---|--|
| Step I  | Identify surveillance activities at each level for each programme/priority disease | Participants             | 1 hour   | Background documents, workshop outputs<br>Flow diagram template | Flow diagram of surveillance structure and process |
| Step II | Analyse tasks at each level for priority diseases                                  | Participants/facilitator | 1 hour   | Task analysis template  | Table of analysed tasks for each priority disease  |

| <b>Step</b> | <b>Specific task</b>   | <b>Person responsible</b> | <b>Duration</b> | <b>Resources</b>        | <b>Output</b>   |
|-------------|--|---------------------------|-----------------|-------------------------|---|
| Step III    | Identify constraints to surveillance at each level and propose realistic solutions | Participants              | 1 hour          | Table of analysed tasks | Table of constraints to surveillance at each level and proposed solutions |

---

# TASK ANALYSIS BY LEVEL FOR PRIORITY DISEASES

(MAY BE PERFORMED AFTER THE ASSESSMENT)

| <b>Task/Activity<br/>Proposed task,<br/>by level</b> | <b>Person<br/>responsible</b> | <b>Timing</b>                  | <b>Skill</b>            | <b>Resources</b>   | <b>Support<br/>function<br/>required</b> |
|--|-------------------------------|--------------------------------|-------------------------|--|--|
| <b>Peripheral Level</b>                              |                               |                                |                         |  |  |
| Detection  | Health worker                 | Per occurrence of health event | Basic diagnostic skills | Written case definitions<br>Register<br>Surveillance forms | Standards<br>Training<br>Supervision     |
| Reporting  |                               |                                |                         |  |  |
| Analysis...  |                               |                                |                         |  |  |
| <b>Intermediate Level</b>                            |                               |                                |                         |  |  |
| Detection/<br>Confirmation                           |                               |                                |                         |  |  |
| Reporting  |                               |                                |                         |  |  |
| Analysis...  |                               |                                |                         |  |  |
| <b>Central Level</b>                                 |                               |                                |                         |  |  |
|  |                               |                                |                         |  |  |



## DESIGNING TOOLS FOR FIELD ASSESSMENT

---

|                      |   |
|----------------------|---|
| Objective:           | To adapt the generic field assessment tools   |
| Method:              | Group discussion  |
| Duration:            | Approximately 8 hours   |
| Materials required:  | List of priority diseases identified and surveillance flow chart  |
| Role of facilitator: | To help the group adapt the generic field assessment questionnaires through group discussions. The facilitator needs to stress the importance of making the generic questions relevant to the country, the need for emphases on pertinent questions, discarding irrelevant ones, regrouping questions, splitting others, and creating new questions if necessary. Although more difficult to analyse, the importance of probing and collecting qualitative data should be stressed. |
| Product:             | Table of performance indicators for surveillance system(s) for the country's priority diseases, field assessment questionnaires for each level (central, district or intermediate, health facility) and laboratory.   |

### **Step I:**

Discuss generic performance indicators and examples with group, then adapt or modify them for the country's priority diseases (through group discussion). Take into account the objectives of surveillance and various components of surveillance that might affect the performance of a system (e.g. available standards, skills, material resources, communication technology).

**Step II:**

Jointly reflect on the various aspects of the surveillance system that need to be assessed at each level (mainly, structure, capacity and synergy within the system, and between systems). Adapt the generic questionnaires (see Annex 12) for field assessment at each level. The questionnaires should be a product of indicators chosen.

---

---

| <b>Step</b> | <b>Specific task</b>  | <b>Person responsible</b>  | <b>Duration</b> | <b>Resources</b>  | <b>Output</b>   |
|-------------|---|----------------------------|-----------------|---|---|
| Step I      | Create/identify indicators to assess system performance for each level for each disease | Facilitators/ participants | 3 hours         | List of priority diseases identified, objective of surveillance | List of indicators to establish the system(s) performance for the priority diseases |
| Step II     | Develop/adapt questionnaires for data collection for indicators at each level           | Facilitators/ participants | 5 hours         | Generic questionnaires  | Questionnaires for field assessment at each level                                   |

---

---

## SELECTION OF ASSESSMENT SITES AND SCHEDULING OF VISITS

|                      |  |
|----------------------|--|
| Objective:           | To select assessment sites, schedule visits and work out logistics   |
| Method:              | Group discussion   |
| Duration:            | Approximately 2 hours  |
| Materials required:  | List of facilities, and maps, template tables  |
| Role of facilitator: | To help the group select assessment sites using acceptable sampling method (see Annex 9.1 for sampling)<br><br>To help the group agree on field visit scheduling and logistics |
| Products:            | Sample sizes for the assessment<br><br>Schedule of field visits and logistic arrangements  |

### Work Plan

| Step     | Specific task   | Person responsible       | Duration   | Resources   | Output   |
|----------|---|--------------------------|------------|---|--|
| Step I   | Selection types of sites and number of facilities to be visited | Participants             | 60 minutes | List of facilities, maps                            | Sample sizes (by level)<br>Table indicating types and number of sites and facilities to be visited, with indication of any exclusions made |
| Step II  | Scheduling field visits   | Participants/facilitator | 30 minutes | Table indicating sites and facilities               | Schedule of field visits for team members  |
| Step III | Arrangement of logistics for field visits                       | Participants/facilitator | 30 minutes | Template table indicating schedules for field visit | Table of equipment, transport, accommodation, security and per-diem arrangements for team members  |



## SELECTION OF SAMPLES FOR REGIONS, DISTRICTS, AND HEALTH FACILITIES

---

The general sampling strategy is to collect information about all levels of the surveillance system; the national, district, health facility levels, including the laboratory. This provides an overall picture of surveillance and response within the health care system.

It may be too expensive and time consuming to use a sample that would enable precise quantitative statements about each characteristic of the surveillance system addressed in the assessment and there may be little added value. Such a sample is not necessarily required, since the purpose of the assessment is to **understand** how the surveillance system is working, in order to address **common problems and challenges, identify synergies and strengthen the system**, rather than to have a scientific statement about the extent of each of the problems. It is particularly important that the sample includes districts representing the broad range of surveillance practices within the country.

One approach to sampling would be to divide the country into a number of strata corresponding to major geographical or administrative areas. Usually administrative regions or provinces have been used.

At the regional or provincial level, each region or province can be further stratified into sub-strata according to important characteristics that affect the functioning of the surveillance system. For example, it might be advantageous to divide the province into areas that appear to have particularly well functioning surveillance systems, those thought to have average systems, and those where it is believed that surveillance is functioning poorly. In addition, if there are areas with particular epidemiological characteristics — such as those prone to certain types of epidemics, where early warning is essential — it might be advisable to include those as separate sub-strata within the region. Districts could be selected randomly within each sub-stratum.

The selection of health facilities requires a detailed list of health facilities, including the level of facility (hospital, health centre or health post) whether they are situated in urban or rural areas, and whether they are public or private. Facilities should then be randomly selected from both rural and urban areas, publicly or privately owned, and representing each type of health facility (hospital, health centre, and health post, dispensaries). Thus, if the district contains rural and urban areas, and public and private health facilities, then health facilities should be selected representing public as well as private facilities in both rural and urban areas.

It is important to keep in mind that the selection of regions or provinces takes place at the national level, while the selection of sample districts takes place at the regional level, and the selection of sample facilities takes place at the district level. There are two reasons for structuring the sampling process in this way. First, one of the main aims of the assessment is to involve all layers of the surveillance system in the process. By selecting the districts and health facilities at the regional and district levels respectively, managers at these levels will feel more involved in the process as a whole. In addition, it is not always the case that the relevant, up-to-date detailed information on districts and their health facilities will be available at the national level.

Sometimes, because of logistic reasons, it will not be possible to visit all parts of the country either because of the remoteness of the area, or because of other reasons that would make visiting the area impossible. These constraints should be identified before the sampling takes place, and the fact that the certain areas were excluded from the sample will need to be taken into consideration in the analysis of the data. If for example, it were not possible to visit any remote areas, this would mean that the sample did not reflect the situation in remote areas, and no conclusions can be drawn about them.

In analyzing the information, it must be remembered that this assessment is not a scientific sample, so that although the data can be summarized, levels of statistical significance cannot be assigned. The analysis should serve to identify common problems in the surveillance system, and suggest areas of the country in which such problems are common.





# CONDUCT DURING FIELD ASSESSMENT

---

## Guide to field communication at different levels

### *Team Members*

1. Introduce team members to each other. This is important to enhance team spirit
2. Identify where, when and how long the assessment will take at each site
3. Explicitly discuss the roles and responsibilities of each team member, which may change from site to site
4. Ensure that the group members have logistics and supply, including data collection tools, stationary, daily allowances, etc.
5. Make sure that there is communication with the overall team leader regularly (daily at the least, recommended)
6. Communicate with the overall co-ordinator before making changes in the tools, field methods or the location. There may be a need to change these. However, changes must be discussed and agreed upon for consistent data collection.

### **Meeting with authorities-focal persons at field**

1. Identify the focal person at the assessment region, zone, facility
2. Plan consultation sessions ahead of time and get it scheduled
3. Introduce team members and brief on mission objectives
4. Outline what your expectations from this briefing meeting are
5. Emphasise that the assessment is for strengthening and making recommendations to facilitate work, and not for critical, judgemental or punitive purposes

6. Invite the focal person to provide views and inputs
7. Agree on roles and accept support from the organizations and institutions supporting surveillance at the field level
8. Explain how you will get feedback of the assessment to them, and any planned follow-up to the mission.

#### **Meeting health workers carrying out surveillance**

1. Give clear description of objectives of the mission
2. Discuss their roles in the assessment (Do they participate and give interviews at lower level? Do they need to be interviewed, have data collected from them, observed executing their practice, etc.)
3. Explain whether you will provide feedback, and if so how it would reach them.

#### **Accessing Communities**

1. Observe and respect community norms
2. Clearly explain the objectives in a simple and concise way.  
Answer their questions
3. Often the mission may raise expectations. Be honest about your mission
4. Select convenient time to conduct community assessments.

## ANALYSIS, PRELIMINARY REPORT WRITING

---

|                      |  |
|----------------------|--|
| Objective:           | To analyse data from field visits and prepare draft report   |
| Method:              | Group discussion   |
| Duration:            | Approximately 3 days   |
| Materials required:  | Products of pre-assessment workshop; questionnaires from the field assessment; data entry and data management skills   |
| Role of facilitator: | To help the groups analyse the data obtained from the field assessment, both qualitative (impressions obtained in the field) and quantitative (questionnaires) and help them draft a preliminary report of the findings  |
| Products:            | Preliminary report of the assessment findings, which will be left in the country assessed<br><br>Draft timetable for writing the final assessment report, for circulation to stakeholders and partners of MoH<br><br>Draft timetable for Plan of Action Workshop |

The preliminary report (see Annex 15 for reporting format) and the draft timetables for writing the final report and the Plan of Action Workshop should be presented at the Post Assessment Workshop.

*Before leaving the country it is important to:*

- Agree on the schedule for follow-up
- Agree on the exact dates for the Plan of Action Workshop
- Arrange for WHO liaison to carry out day to day follow-up with MoH focal point regarding the preparation and circulation of the final assessment report
- Organize regular updates on progress and involve technical and donor partners within the country
- WHO should be informed about any major obstacles encountered.



# PROTOTYPE REPORT WRITING FORMAT

---

## **Executive Summary**

### **1. Introduction**

### **2. Background on the country**

#### 2.1 Geography

#### 2.2 Demography

#### 2.3 Socio-economic factors

#### 2.4 Health systems

##### 2.4.1 Health services infrastructure

##### 2.4.2 Human resources for health

##### 2.4.3 Health status (description, indicators)

##### 2.4.4 The burden of disease (mortality, morbidity, infectious diseases)

##### 2.4.5 Decentralization (if relevant)

##### 2.4.6 The health sector strategic plan if relevant

##### 2.4.7 Review of existing surveillance systems (include flow chart, organogramme)

##### 2.4.8 Brief description of existing components of systems assessed

###### 2.4.8.1 Priority diseases

###### 2.4.8.2 Structure

###### 2.4.8.3 Process/Capacity

###### 2.4.8.4 Out put

###### 2.4.8.5 Integration

###### 2.4.8.6 Laboratories

###### 2.4.8.7 GIS

###### 2.4.8.8 Communication

### **3. Objectives of assessment**

- 3.1 General objective
- 3.1 Specific objectives

### **4. Methodology**

- 4.1 Preparation for the assessment
- 4.2 Selection of sites
  - 4.2.1 Selection of regions/provinces
  - 4.2.2 Selection of districts.
  - 4.2.3 Selection of health facilities
- 4.3 Procedure and data collection tools
- 4.4 Composition of assessment teams
- 4.5 Training of assessment teams
- 4.6 Field testing
- 4.7 Field assessment
- 4.8 Data analysis
  - 4.8.1 Quantitative analysis
  - 4.8.2 Qualitative analysis

### **5. Findings: For each level**

- 5.1.1 Presence of surveillance systems
- 5.1.2 Availability of case definition (health facility)
- 5.1.3 Case confirmation (health facility)
- 5.1.4 Data reporting (completeness and timeliness)
- 5.1.5 Data analysis
- 5.1.6 Outbreak investigation
- 5.1.7 Epidemic preparedness
- 5.1.8 Epidemic response
- 5.1.9 Feedback
- 5.1.10 Supervision
- 5.1.11 Co-ordination
- 5.1.12 Training
- 5.1.13 Resources
- 5.1.14 The laboratory
- 5.1.15 GIS
- 5.1.16 Communications

**6. Conclusion**

**7. Recommendations**

**8. Annexes**

Example: Annex 1. Qualitative analysis (strengths, weaknesses, opportunities, threats, solutions/recommendations)

- 8.1 Existence of the surveillance systems
- 8.2 Case detection
- 8.3 Case registration
- 8.4 Case confirmation
- 8.5 Reporting
- 8.6 Feedback from higher levels
- 8.7 Data analysis
- 8.8 Epidemic preparedness and response
- 8.9 Training
- 8.10 Supervision
- 8.11 Surveillance co-ordination
- 8.12 Resources
- 8.13 Conclusions and recommendations



## POA MATRIX

**Building on the findings and recommendations from the assessment**

|                       | Goals | Objectives | Activities | Timeline | Implementers | Resources | Obstacles | Indicators |
|-----------------------|-------|------------|------------|----------|--------------|-----------|-----------|------------|
| Case detection        |       |            |            |          |              |           |           |            |
| Registration          |       |            |            |          |              |           |           |            |
| Confirmation          |       |            |            |          |              |           |           |            |
| Reporting             |       |            |            |          |              |           |           |            |
| Analysis              |       |            |            |          |              |           |           |            |
| Response              |       |            |            |          |              |           |           |            |
| Epidemic preparedness |       |            |            |          |              |           |           |            |
| Communication         |       |            |            |          |              |           |           |            |
| Training              |       |            |            |          |              |           |           |            |
| Supervision           |       |            |            |          |              |           |           |            |
| Feedback              |       |            |            |          |              |           |           |            |
| Laboratory            |       |            |            |          |              |           |           |            |
| Integration           |       |            |            |          |              |           |           |            |

