


Estimated number of people needing antiretroviral therapy (0-49 years), 2005: **1 350**  
 Antiretroviral therapy target declared by country: **700 by the end of 2005**



Map Data Source:  
 WHO/UNAIDS Epidemiological Fact Sheets  
 and the United States Census Bureau  
 Map production:  
 Public Health Mapping & GIS  
 Communicable Diseases (CDS)  
 World Health Organization



## 1. Demographic and socioeconomic data

	Date	Estimate	Source
Total population (millions)	2004	0.7	United Nations
Population in urban areas (%)	2005	84.568	United Nations
Life expectancy at birth (years)	2003	55	WHO
Gross domestic product per capita (US\$)	2002	855	Banque Nationale de Djibouti
Government budget spent on health care (%)	2002	10.1	WHO
Per capita expenditure on health (US\$)	2002	54	WHO
Human Development Index	2003	0.495	UNDP

°= Percentage of young people 15-24 years who correctly identify two major ways of preventing the sexual transmission of HIV (using condoms and limiting sex to one faithful, uninfected partner), who reject the two most common local misconceptions about HIV transmission, and who know that a healthy looking person can transmit HIV.

°°=Percentage of young people 15-24 years reporting the use of a condom during sex with a non-regular partner in the last 12 months.

one site only

## 2. HIV indicators

	Date	Estimate	Source
Adult prevalence of HIV/AIDS (15-49 years)	2003	0.7% - 7.5%	WHO/UNAIDS
Estimated number of people living with HIV/AIDS (0-49 years)	2003	2 300 - 24 000	WHO/UNAIDS
Reported number of people receiving antiretroviral therapy (0-49 years), 2005	Sep 2005	356	TGF
Estimated number of people needing antiretroviral therapy (0-49 years), 2005	Dec 2004	1 350	WHO/UNAIDS
HIV testing and counselling sites: number of sites	Oct 2005	8	NAP
HIV testing and counselling sites: number of people tested at all sites	1Jan - 30 Sep 05	1,930*	NAP
Knowledge of HIV prevention methods (15-24 years)% - female°			
Knowledge of HIV prevention methods (15-24 years)% - male°			
Reported condom use at last higher risk sex (15-24 years)% - female°°			
Reported condom use at last higher risk sex (15-24 years)% - male°°			

## 3. Situation analysis

Epidemic level and trend and gender data

Major vulnerable and affected groups

Policy on HIV testing and treatment

Antiretroviral therapy: first-line drug regimen, cost per person per year

Assessment of overall health sector response and capacity

Critical issues and major challenges

## 4. Resource requirements and funds committed for scaling up treatment and prevention in 2004-2005



## 5. Treatment and prevention coverage

## 6. Implementation partners involved in scaling up treatment and prevention

Leadership and management

Service delivery

Community mobilization

Strategic information

## 7. Staffing input for scaling up HIV treatment and prevention

WHO's response so far

Key areas for WHO support in the future

Staffing input for scaling up HIV treatment and prevention