










**Figure 1 Global number of adults (15+ years) eligible for ART in low- and middle-income countries, by 2006 and 2010 WHO Guidelines, 1990-2009**

- Adults (15+ years) eligible for ART in low and middle-income countries; CD4 count < 350
- High and low estimates
- Adults (15+ years) eligible for ART in low and middle-income countries; CD4 count < 200
- High and low estimates




**Figure 2** Number of adults (15+ years) eligible for ART in **low- and middle-income countries**, by region and by 2006 and 2010 WHO Guidelines, 1990-2009




**Figure 3** Number of adults (15+ years) eligible for antiretroviral therapy in Sub-Saharan Africa, 2009

2010 WHO recommendations: **CD4 < 350**



2006 WHO recommendations: **CD4 < 200**



**Table 1** Difference in the number eligible for antiretroviral therapy (ART) in 2009 in sub-Saharan Africa: 2006 WHO guidelines compared with 2010 WHO guidelines

Country	ART need in 2009			Proportional increase between the two guidelines	Coverage of ART based on 2006 guidelines (%)	Number of people on ART
	Based on 2006 guidelines	Based on 2010 guidelines	Difference*			
Botswana	140 000	170 000	33 000	0.24	103	150 000
Cameroon	180 000	270 000	85 000	0.46	41	76 000
Central African Republic	51 000	74 000	23 000	0.45	28	14 000
Kenya	520 000	700 000	190 000	0.37	65	340 000
Lesotho	90 000	130 000	39 000	0.43	68	62 000
Malawi	310 000	440 000	120 000	0.39	63	200 000
Mozambique	380 000	570 000	190 000	0.51	39	170 000
South Africa	1700 000	2600 000	900 000	0.51	56	970 000
Swaziland	56 000	80 000	24 000	0.43	85	47 000
Uganda	380 000	520 000	140 000	0.36	54	200 000
United Republic of Tanzania	450 000	660 000	210 000	0.45	44	200 000
Zambia	330 000	440 000	110 000	0.33	85	280 000
Zimbabwe	450 000	640 000	190 000	0.43	49	220 000