



Global Immunization Data

November 2011

Summary: Global immunization coverage in 2010

Based on the latest World Health Organization (WHO)/UNICEF global estimates for 2010, trends related to global vaccination coverage continue to be positive.

Immunization currently averts an estimated 2.5 million deaths every year in all age groups from diphtheria, tetanus, pertussis (whooping cough), and measles.

More children are being reached with immunization. In 2010, an estimated 109 million children under the age of one were vaccinated with three doses of diphtheria-tetanus-pertussis (DTP3) vaccine.

More countries achieve high levels of vaccination coverage

Three regions — the Americas, Europe and Western Pacific — maintained over 90% immunization coverage.

Number of countries reaching 90% or more immunization coverage with DTP3 vaccine in 2010: 130 countries compared to 123 in 2009.

However the number of countries reaching over 80% DTP3 coverage remained at 157 in 2009 and 2010.

Increasing uptake of new and underused vaccines

Hepatitis B vaccine for infants was introduced nationwide in 179 countries (including in parts of India and the Sudan) by the end of 2010. Global hepatitis B vaccine coverage is estimated at 75% and is as high as 91% in the Western Pacific and 89% in the Americas. Coverage in the South-East Asia Region reached 52% in 2010.

Haemophilus influenzae type B (Hib) vaccine was introduced in 169 countries by the end of 2010 (including in parts of Belarus, the Philippines and the Sudan), up from 161 countries in 2009. Global coverage with three doses of Hib vaccine is estimated at 42% in 2010, reaching 92% in the Americas, but only 9% and 10% in the South-East Asia Region and in the Western Pacific Region, respectively.

Rubella vaccine was introduced nationwide in 131 countries by the end of 2010, up from 83 countries in 1996. There has been remarkable progress towards the elimination of rubella and congenital rubella syndrome in the Americas with a reduction of 99.99% of confirmed cases between 1998 and 2010.

Mumps vaccine was introduced nationwide in 118 countries by the end of 2010.

Yellow fever vaccine was introduced in routine infant immunization programmes in 36 countries and territories out of the 47 at risk for yellow fever in Africa and the Americas.

Maternal and neonatal tetanus (MNT): the vaccine to prevent MNT was introduced as part of routine immunization programmes in over 100 countries by the end of 2010. Vaccination coverage with at least two doses of tetanus toxoid vaccine or tetanus-diphtheria toxoid vaccine was estimated at 68% in 2010 and an estimated 84% of newborns were protected against neonatal tetanus through immunization. As of December 2010, maternal and neonatal tetanus persist as public health problems in 38 countries, mainly in Africa and Asia.

Pneumococcal vaccine was introduced in 55 countries (including one country where the vaccine was partially introduced) by the end of 2010, up from 44 countries in 2009.

Rotavirus vaccine was introduced in 28 countries by the end of 2010, up from 23 in 2009.

Human papillomavirus (HPV) vaccine was introduced in 37 countries (including four in parts of the country only) by the end of 2010, up from 29 countries by the end of 2009.

The unprotected children

Number of children under one year of age who did not receive DTP3 vaccine worldwide: 19.3 million in 2010 compared to 19.7 million in 2009.

Nearly seventy percent of these children live in ten countries: Afghanistan, Democratic Republic of the Congo, Ethiopia, India, Indonesia, Iraq, Nigeria, Pakistan, South Africa and Uganda.

Deaths due to vaccine-preventable diseases

Total number of children who died from diseases preventable by vaccines currently recommended by WHO: 1.7 million^a.

- Hib: 260 000^b
- Pertussis: 195 000^a
- Measles: 118 000^a
- Neonatal tetanus: 59 000^a
- Tetanus (non-neonatal): 2 000^a
- Pneumococcal disease: 520 000^b
- Rotavirus: 527 000^b

Estimated number of all deaths in children under five in 2008: 8.8 million.
Nearly 20% of all deaths in children under five is vaccine preventable

Estimated number of all deaths in children 1-59 months of age: 5.2 million.
30% of deaths in children 1-59 months of age are vaccine preventable.

DTP

Global coverage of infants in 2010 with DTP3 vaccine: 85%^c.

Global coverage of infants in 1990 with DTP3 vaccine: 75%^c.

Estimated number of children vaccinated with DTP3 vaccine in 2010: 109 million^c.

Polio

Global coverage of infants with three doses of polio vaccine in 2010: 86^c.

Global coverage of infants with three doses of polio vaccine in 1990: 75%^c.

Reported number of polio cases in 2010: 1410 confirmed polio cases (including 1349 wild virus confirmed cases)^d.

Estimated number of polio cases in 1988: 350 000.

Number of polio-endemic countries in 2010: 4.

Number of polio-endemic countries in 1988: 125.

Measles

Global coverage of children by their second birthday with one dose of measles-containing vaccine in 2010: 85%^c.

Global coverage of children by their second birthday with one dose of measles-containing vaccine in 1990: 73%^c.

Number of countries with a second dose of measles vaccine in routine immunization schedule: 136 (70% of 193 countries)^e.

Number of estimated measles deaths in 2008: 164 000 [115 000 - 222 000]*

MNT

Number of countries that had not yet eliminated MNT in 2010: 38^f.

Number of countries that had not yet eliminated MNT in 2000: 58^f.

Number of women living in high-risk areas protected with at least two doses of tetanus toxoid vaccine given during supplementary immunization activities (1999-2010): more than 100 million^f.

Hepatitis B

Global coverage of infants with three doses of hepatitis B vaccine in 2010: 75%^c.

Global coverage of infants with three doses of hepatitis B vaccine in 1990: 1%^c.

Yellow Fever

Global yellow fever vaccine coverage in 2010: 43% (includes only the 47 countries and territories at risk)^e.

Notes:

All numbers referring to deaths are rounded to the nearest thousand (000).

WHO has 193 Member States.

*Estimates of the uncertainty intervals have been made by assuming a probability distribution around the immunization coverage, vaccine efficacy, age-specific distributions of cases, and age-specific case-fatality rates. From these distributions 10 000 simultaneous samples of the parameter values were drawn and the estimate of deaths re-calculated. The uncertainty intervals represent the interval within which 95% of all 10 000 sample mortality estimates occurred.

More information, including more detailed immunization data and graphs, can be found at http://www.who.int/immunization_monitoring/data/en/.

^a Global Burden of disease 2008 update, WHO.

^b WHO/IVB provisional estimates based on Global Burden of Diseases estimates.

^c WHO/UNICEF immunization coverage estimates, data as of July 2011.

^d Polio Eradication programme data as of September 2011.

^e WHO Department of Immunization, Vaccines and Biologicals, data as of September 2011.

^f WHO/UNICEF MNT data as of September 2011.