

BOTSWANA

Last Updated: 2008-01-21

Level	Date	Region and sample descriptor	Sex	Age (years)	Sample size	Haemoglobin (g/L)						Reference	Notes				
						Proportion (%) of population with haemoglobin below:							Mean	SD	Method	General	Line
						70	100	110	115	120	130						
N	1994	NPW: Total	F	15.00- 49.99	315	1.3					32.7		17	C	2805	*	
		Pre-SAC by age	B	0.50- 4.99	149	0.0		38.0						14			
		SAC by age	B	5.00- 8.99	100	0.0			53.0					14			
L	1971	3 villages	B	0.00- 5.99	754			22.0						D	2265	*	1
		By village: Kumakwane	B	0.00- 5.99	256			18.0									
		By village: Good Hope	B	0.00- 5.99	415			25.0									
		By village: Khudumelapye	B	0.00- 5.99	83			23.0									

NOTES

BOTSWANA

Reference No: 2805

General Notes: *Data from raw data not report.*

Reference No: 2265

General Notes: *Method: Spencer haemoglobin meter.*

Line note 1 Disaggregated data by age.

REFERENCES

BOTSWANA

Reference 2265 Burgees JL. Nutrition Survey in Botswana. June-July 1971. 1971.

Reference 2805 Ministry of Health [Botswana], UNICEF. Micronutrient malnutrition in Botswana. A national survey to assess the status of iodine, iron, and vitamin A. Gaborone, Ministry of Health, 1996.

ADDITIONAL REFERENCES

BOTSWANA

- Reference 782 Ministry of Health. A review paper for the food and nutrition situation in Botswana. Rome, 1992.
- Reference 2343 WHO Regional Office for Africa. Anaemia in the African region. Epidemiological Bulletin, 1989, 7 :20-37.
- Reference 5067 Abrams SA, Mushi A, Hilmers DC, Griffin IJ, Davila P, Allen L. A multivitamin-fortified beverage enhances the nutritional status of children in Botswana. Journal of Nutrition, 2003, 133 :1834-1840.