

Health service coverage

56835
83347-9473

Figures have been computed by WHO to ensure comparability; thus they are not necessarily the official statistics of Member States, which may use alternative rigorous methods.

	Member State	WHO region	Immunization coverage among 1-year-olds ^a			Antenatal care coverage ^b			Births attended by skilled health personnel ^c		Contraceptive prevalence rate ^d	
			Measles	DTP3	HepB3	At least 1 visit	At least 4 visits	Year				
			(%) 2005	(%) 2005	(%) 2005	(%)	(%)		(%)	Year	(%)	Year
1	Afghanistan	EMR	64	76	...	52	...	2003	14	2003	4.8	2000
2	Albania	EUR	97	98	98	81	42	2002	94	2002 ^k	75.1	2002
3	Algeria	AFR	83	88	83	79	...	2000	90	2002	64.0	2000
4	Andorra	EUR	94	98	79	
5	Angola	AFR	45	47		45	2001	6.2	2001
6	Antigua and Barbuda	AMR	99	99	99	...	82	2001	100	2005	...	
7	Argentina	AMR	99	92	87	...	95	2001	99	2004 ^k	...	
8	Armenia	EUR	94	90	91	82	65	2000	98	2005	60.5	2000
9	Australia	WPR	94	92	94		100	2003	...	
10	Austria	EUR	75	86	86	
11	Azerbaijan	EUR	98	93	96	70	...	2001	74	2001	55.4	2001
12	Bahamas	AMR	85	93	93		99	2004	...	
13	Bahrain	EMR	99	98	98		99	2005	...	
14	Bangladesh	SEAR	81	88	62	39	11	1999–2000	13	2004 ^k	58.1	2004
15	Barbados	AMR	93	92	92	89	...	2001	100	2004	...	
16	Belarus	EUR	99	99	99		100	2005	...	
17	Belgium	EUR	88	97	78		99	1999	...	
18	Belize	AMR	95	96	97	...	96	2001	89	2005	...	
19	Benin	AFR	85	93	92	88	61	2001	66	2001 ^k	18.6	2001
20	Bhutan	SEAR	93	95	95		51	2005	...	
21	Bolivia	AMR	64	81	81	84	69	2001	61	2003–2004 ^k	58.4	2003–2004
22	Bosnia and Herzegovina	EUR	90	93	93	99	...	2000	100	2000	47.5	2000
23	Botswana	AFR	90	97	85	99	97	2001	99	2000	40.4	2000
24	Brazil	AMR	99	96	92		97	2003	...	
25	Brunei Darussalam	WPR	97	99	99	...	100	2001	100	2004	...	
26	Bulgaria	EUR	96	96	96		99	2005	41.5	1997
27	Burkina Faso	AFR	84	96	...	72	18	2003	38	2003	13.8	2003
28	Burundi	AFR	75	74	74	93	79	2001	25	2000	15.7	2000
29	Cambodia	WPR	79	82	...	44	9	2000	44	2005–2006	23.8	2000
30	Cameroon	AFR	68	80	79	77	52	1998	62	2004	26.0	2004
31	Canada	AMR	94	94		100	2003–2004	...	
32	Cape Verde	AFR	65	73	69	...	99	2001	89	1998 ^k	52.9	1998
33	Central African Republic	AFR	35	40		44	2000	27.9	2000
34	Chad	AFR	23	20	...	51	13	1997	15	2004–2005 ^k	2.8	2004
35	Chile	AMR	90	91		100	2004	...	
36	China	WPR	86	87	84		83	2004	90.2	2004
37	Colombia	AMR	89	87	87	90	79	2000	91	2004–2005	78.2	2004–2005
38	Comoros	AFR	80	80	80		62	2000	25.7	2000
39	Congo	AFR	56	65		83	2005–2006 ^k	44.3	2005
40	Cook Islands	WPR	99	99	99		98	2001	...	
41	Costa Rica	AMR	89	91	90	...	70	2001	98	2004	80.0	1999
42	Côte d'Ivoire	AFR	51	56	56	84	35	1998–1999	63	2000	15.0	1998–1999
43	Croatia	EUR	96	96	99		100	2005	...	
44	Cuba	AMR	98	99	99	...	100	2001	100	2005	73.3	2000
45	Cyprus	EUR	86	98	88		99	2003	...	



Children aged <5 years sleeping under insecticide-treated bednets ^e		Antiretroviral therapy coverage		TB detection rate under DOTS ^h	TB treatment success under DOTS ⁱ	Children aged < 5 years with ARI symptoms taken to facility ^j		Children aged < 5 years with diarrhoea receiving ORT ^l		Children aged < 5 years with fever who received treatment with any antimalarial ^o		Children 6–59 months who received vitamin A supplementation ^l		Births by Caesarean section ^b	
(%)	Year	(%) Dec 2006	(%) Dec 2005			(%) 2005	(%) 2004 cohort	(%)	Year	(%)	Year	(%)	Year	(%)	Year
...	44	89
...	25	78	15	2002
...	...	25	...	106	91	6	2000
...	94	100	24	1999
2	2001	10	1	85	68	63.0	2001
...	246	100
...	...	79	87	67	58
...	60	70	24.6	2000	59.7	2000	7	2000
...	42	85	21	1998
...	56	69	21	2002
1	2000	<1	...	55	60	0.8	2000	4	2002
...
...	77	82
...	...	<3	...	59	90	20.3	2004	83.4	2004	81.8	2004	3	1999–2000
...	...	87
...	...	20	...	46	74	17	2002
...	64	72	16	1999
...	...	59	...	102	60
7	2001	38	17	83	83	29.3	2001	60.9	2001	60.4	2001	4	2001
...	31	83
...	...	24	...	72	80	51.5	2003	66.4	2003	60.0	2003	15	1998
...	71	98
...	...	>95	54	69	65
...	...	85	48	53	81
...	112	71
...	90	80	17	2002
2	2003	39	6	18	67	32.6	2003	62.8	2003	49.6	2003	33.3	2003	1	2003
1	2000	26	3	30	78	31.3	2000
...	4	66	91	34.5	2000	74.1	2000	28.5	2000	1	2000
1	2004	25	10	106	71	39.9	2004	56.7	2004	53.1	2004	37.5	2004	3	1998
...	64	62	19	1997–1998
...	34	71	6	1998
2	2000	6	4	40	91	68.8	2000
1	2000	14	1	22	69	11.8	2004	37.5	2004	55.8	2004	32.0	2004	1	1996–1997
...	...	83	...	112	83
...	...	27	2	80	94
3	2000	50	...	26	85	56.7	2005	70.1	2005	25	2000
9	2000	49	94	62.7	2000
...	...	17	10	57	63	43.9	2005	53.5	2005	65.5	2005
...	77	0
...	...	>95	...	118
1	2000	28	4	38	71	34.9	1999	66.1	1999	57.5	2000	3	1998–1999
...	14	2002
...	...	>95	...	98	93
...	57	20

Health service coverage

56835
83347-9473

Figures have been computed by WHO to ensure comparability; thus they are not necessarily the official statistics of Member States, which may use alternative rigorous methods.

	Member State	WHO region	Immunization coverage among 1-year-olds ^a			Antenatal care coverage ^b			Births attended by skilled health personnel ^c		Contraceptive prevalence rate ^d	
			Measles	DTP3	HepB3	At least 1 visit	At least 4 visits	Year	Year	Year	Year	
			(%) 2005	(%) 2005	(%) 2005	(%)	(%)	(%)				
46	Czech Republic	EUR	97	97	99	100	2005	72.0	1997
47	Democratic People's Republic of Korea	SEAR	96	79	92	98	...	2000	97	2004
48	Democratic Republic of the Congo	AFR	70	73	61	2001	31.4	2001
49	Denmark	EUR	95	93
50	Djibouti	EMR	65	71	61	2003
51	Dominica	AMR	98	98	100	2001	100	2004
52	Dominican Republic	AMR	99	77	77	100	93	1999	98	2002 ^a	69.8	2002
53	Ecuador	AMR	93	94	94	56	...	1999	74	2004	65.8	1999
54	Egypt	EMR	98	98	98	54	41	2000	74	2005	59.2	2005
55	El Salvador	AMR	99	89	89	...	76	2001	69	2002–2003	67.3	2002–2003
56	Equatorial Guinea	AFR	51	33	37	2001	65	2000
57	Eritrea	AFR	84	83	83	...	49	2001	28	2002	8.0	2002
58	Estonia	EUR	96	96	95	100	2005
59	Ethiopia	AFR	59	69	...	27	10	2000	6	2005	14.7	2005
60	Fiji	WPR	70	75	75	99	2002
61	Finland	EUR	97	97	100	2002
62	France	EUR	87	98	29
63	Gabon	AFR	55	38	55	94	63	2000	86	2000 ^a	32.7	2000
64	Gambia	AFR	84	88	88	92	...	2000	55	2000	9.6	2000
65	Georgia	EUR	92	84	74	91	...	1999	92	2005	47.2	2005
66	Germany	EUR	93	90	84	100	2006
67	Ghana	AFR	83	84	84	90	69	2003	47	2003	25.2	2003
68	Greece	EUR	88	88	88
69	Grenada	AMR	99	99	99	...	98	2001	100	2005
70	Guatemala	AMR	77	81	27	86	68	1998–1999	41	2002 ^a	43.3	2002
71	Guinea	AFR	59	69	...	74	48	1999	38	2005	9.1	2005
72	Guinea-Bissau	AFR	80	80	...	89	62	2001	35	2000	7.6	2000
73	Guyana	AMR	92	93	93	88	...	2000	86	2000 ^a	37.3	2000
74	Haiti	AMR	54	43	...	79	42	2000	25	2005–2006	28.1	2000
75	Honduras	AMR	92	91	91	...	84	2001	67	2005–2006	61.8	2001
76	Hungary	EUR	99	99	100	2005
77	Iceland	EUR	90	95
78	India	SEAR	58	59	8	65	30	1998–1999	48	2005–2006 ^a	48.2	1998–1999
79	Indonesia	SEAR	72	70	70	97	81	2002–2003	66	2002–2003	60.3	2002–2003
80	Iran (Islamic Republic of)	EMR	94	95	94	...	77	2001	90	2000	72.9	1997
81	Iraq	EMR	90	81	81	...	78	2001	72	2000
82	Ireland	EUR	84	90	100	2002
83	Israel	EUR	95	95	95
84	Italy	EUR	87	96	96	99	2003
85	Jamaica	AMR	84	88	87	...	99	2001	97	2002–2003	65.9	1997
86	Japan	WPR	99	99	100	2004	55.9	2000
87	Jordan	EMR	95	95	95	99	91	2002	100	2002	55.8	2002
88	Kazakhstan	EUR	99	98	94	82	71	1999	99	2005	66.1	1999
89	Kenya	AFR	69	76	76	88	52	2003	42	2003	39.3	2003
90	Kiribati	WPR	56	62	67	...	88	2001	89	2002



Children aged <5 years sleeping under insecticide-treated bednets ^e		Antiretroviral therapy coverage		TB detection rate under DOTS ^h	TB treatment success under DOTS ⁱ	Children aged < 5 years with ARI symptoms taken to facility ^j		Children aged < 5 years with diarrhoea receiving ORT ^l		Children aged < 5 years with fever who received treatment with any antimalarial ^o		Children 6–59 months who received vitamin A supplementation ^l		Births by Caesarean section ^b	
(%)	Year	(%) Dec 2006	(%) Dec 2005			(%) 2005	(%) 2004 cohort	(%)	Year	(%)	Year	(%)	Year	(%)	Year
...	65	73	14	2002
...	99	89
1	2001	11	2	72	85	45.4	2001
...	71	88	18	2001
...	...	20	1	42	80
...
...	...	37	27	76	80	60.8	2002	55.0	2002	30.7	2002	32	1999
...	...	34	22	28	85	19	1999
...	...	22	...	63	70	72.6	2005	35.7	2005	13.8	2005	11	2000
...	...	39	18	67	90	16	1998
1	2000	24	48.6	2000
4	2002	8	2	13	85	43.6	2002	68.4	2002	3.6	2002	38.0	2002
...	64	71	15	2002
...	33	79	18.7	2005	37.1	2005	3.0	2000	45.8	2005	1	2000
...	72
...	16	2002
...	16	1999
...	...	35	3	57	40	47.7	2000	71.6	2000	6	2000
15	2000	12	6	69	86	55.2	2000
...	...	32	...	91	68	12	2002
...	52	68	22	2001
4	2003	16	6	37	72	44.0	2003	63.3	2003	62.8	2003	78.4	2003	4	2003
...
...
1	2000	52	10	55	85	37.4	1999	58.6	1999	12	1998–1999
...	...	10	1	56	72	34.5	2005	56.7	2005	68.2	2005	2	1999
7	2000	6	...	79	75	58.4	2000
...	...	72	...	40	72	2.6	2000
...	...	39	7	57	80	36.3	2000	54.9	2000	11.7	2000	31.6	2000	2	2000
...	...	40	12	82	85	55.9	2005	66.8	2005
...	43	54	23	2002
...	53	50	17	2001
...	2	61	86	7	1998–1999
0	2000	20	<1	66	90	56.8	2003	60.6	2003	0.7	2002–2003	75.1	2003	4	2002–2003
...	...	5	...	64	84
<0.1	2000	0	...	43	85
...	19	2000
...	42	80	17	2001
...	72	32	1999
...	...	56	...	61	46
...	57	57
...	63	85	71.7	2002	63.9	2002	16	2002
...	...	10	...	72	72	48.0	1999	52.6	1999	11	1998
5	2003	44	20	43	80	45.5	2003	50.6	2003	26.5	2003	33.3	2003	4	2003
...	73	94

Health service coverage

56835
83347-9473

Figures have been computed by WHO to ensure comparability; thus they are not necessarily the official statistics of Member States, which may use alternative rigorous methods.

	Member State	WHO region	Immunization coverage among 1-year-olds ^a			Antenatal care coverage ^b			Births attended by skilled health personnel ^c		Contraceptive prevalence rate ^d	
			Measles	DTP3	HepB3	At least 1 visit	At least 4 visits	Year	Year	Year	Year	
			(%) 2005	(%) 2005	(%) 2005	(%)	(%)	(%)				
91	Kuwait	EMR	99	99	99	100	2002
92	Kyrgyzstan	EUR	99	98	97	88	81	1997	98	2005	59.5	1997
93	Lao People's Democratic Republic	WPR	41	49	49	44	29	2001	19	2001	32.2	2000
94	Latvia	EUR	95	99	98	100	2005
95	Lebanon	EMR	96	92	88	...	87	2001	98	2004
96	Lesotho	AFR	85	83	83	91	88	2001	55	2004 ^a	37.3	2004
97	Liberia	AFR	94	87	84	2001	51	1999–2000
98	Libyan Arab Jamahiriya	EMR	97	98	97	...	81	2001	99	1999
99	Lithuania	EUR	97	94	95	100	2005
100	Luxembourg	EUR	95	99	95	100	2002
101	Madagascar	AFR	59	61	61	91	38	1997	51	2003–2004	27.1	2003–2004
102	Malawi	AFR	82	93	93	94	55	2000	56	2004–2005	32.5	2004
103	Malaysia	WPR	90	90	90	100	2005
104	Maldives	SEAR	97	98	98	98	81	2001	70	2001 ^a	42.0	1999
105	Mali	AFR	86	85	85	53	30	2001	41	2001	8.1	2001
106	Malta	EUR	86	92	78	100	2006
107	Marshall Islands	WPR	86	77	89	95	2002
108	Mauritania	AFR	61	71	42	63	16	2000–2001	57	2000–2001 ^a	8.0	2000–2001
109	Mauritius	AFR	98	97	97	99	2005	75.9	2002
110	Mexico	AMR	96	98	98	...	86	2001	93	2003	68.4	1997
111	Micronesia (Federated States of)	WPR	96	94	91	88	2001
112	Monaco	EUR	99	99	99
113	Mongolia	WPR	99	99	98	...	97	2001	100	2004	67.4	2000
114	Montenegro	EUR
115	Morocco	EMR	97	98	96	63	2003–2004	63.0	2003–2004
116	Mozambique	AFR	77	72	72	71	41	1997	48	2003–2004	16.5	2003–2004
117	Myanmar	SEAR	72	73	62	...	76	2001	68	2003	37.0	2001
118	Namibia	AFR	73	86	...	85	69	2000	76	2000	43.7	2000
119	Nauru	WPR	80	80	80
120	Nepal	SEAR	74	75	41	49	15	2001	19	2006	39.3	2001
121	Netherlands	EUR	96	98	100	2003
122	New Zealand	WPR	82	89	87	97	2001
123	Nicaragua	AMR	96	86	86	85	72	2001	67	2001	68.6	2001
124	Niger	AFR	83	89	...	39	11	1998	16	2000 ^a	14.0	2000
125	Nigeria	AFR	35	25	...	61	47	2003	35	2003 ^a	12.6	2003
126	Niue	WPR	99	85	86	100	2005
127	Norway	EUR	90	91
128	Oman	EMR	98	99	99	98	2005
129	Pakistan	EMR	78	72	73	31	2004–2005	27.6	2000–2001
130	Palau	WPR	98	98	98	100	2002
131	Panama	AMR	99	85	85	...	72	2001	91	2004
132	Papua New Guinea	WPR	60	61	63	...	78	2001	42	2004
133	Paraguay	AMR	90	75	75	...	89	2001	77	2004	72.8	2004
134	Peru	AMR	80	84	84	85	69	2000	71	2004 ^a	70.5	2003–2004
135	Philippines	WPR	80	79	44	94	70	2003	60	2003	48.9	2003



Children aged <5 years sleeping under insecticide-treated bednets ^e		Antiretroviral therapy coverage		TB detection rate under DOTS ^h	TB treatment success under DOTS ⁱ	Children aged < 5 years with ARI symptoms taken to facility ^j		Children aged < 5 years with diarrhoea receiving ORT ^k		Children aged < 5 years with fever who received treatment with any antimalarial ^g		Children 6–59 months who received vitamin A supplementation ^l		Births by Caesarean section ^b	
(%)	Year	(%) Dec 2006	(%) Dec 2005			(%) 2005	(%) 2004 cohort	(%)	Year	(%)	Year	(%)	Year	(%)	Year
...	66	63
...	67	85	73.5	1997	6	1997
15	2002	68	86
...	83	73	17	2002
...	74	90	23	1998
...	...	31	12	85	69	54.4	2005	79.6	2005	54.6	2005
...	50	70
...	178	64
...	100	72	15	2002
...	59	19	2000
0	2000	1	<1	67	71	39.3	2004	58.1	2004	41.1	2004	76.2	2004	1	1997
36	2004	43	6	39	71	19.6	2004	70.1	2004	31.6	2004	65.4	2004	3	2000
...	...	22	10	73	56
...	94	95
...	...	37	3	21	71	42.8	2001	65.7	2001	37.6	2003	40.9	2001	1	2001
...	50	100	25	2002
...	77	90
2	2003–2004	17	1	28	22	39.4	2001	47.7	2001	33.4	2003–2004	57.8	2001	3	2000–2001
...	32	89
...	...	76	...	110	82
...	61	80
...
...	82	88	5	2000
...
...	...	41	...	101	87	34.5	2004	54.0	2004	25.5	2004
...	...	14	6	49	77	51.4	2003	70.5	2003	49.8	2003	3	1997
...	...	7	8	95	84
...	...	71	29	90	68	53.1	2000	65.8	2000	14.4	2000	38.1	2000
...
...	...	4	...	67	87	23.7	2001	46.5	2001	81.0	2001	1	2001
...	47	83	14	2002
...	51	66	19	1999
...	...	35	...	88	87	57.4	2001	67.7	2001	1.8	2001	65.3	2001	15	2001
1	2000	6	1	50	61	28.1	2006	52.9	2006	48.1	2000	1	1998
1	2003	15	<1	22	73	31.4	2003	40.2	2003	33.8	2003	33.7	2003	2	2003
...	0	0
...	44	89	16	2001
...	108	90
...	...	<1	...	37	82
...	64	100
...	...	70	...	131	78
...	...	8	1	21	65
...	...	64	...	33	83
...	...	50	9	86	90	68.0	2004	70.6	2004	13	2000
...	...	10	...	75	87	46.3	2003	58.9	2003	76.0	2003	7	2003

Health service coverage

56835
83347-9473

Figures have been computed by WHO to ensure comparability; thus they are not necessarily the official statistics of Member States, which may use alternative rigorous methods.

	Member State	WHO region	Immunization coverage among 1-year-olds ^a			Antenatal care coverage ^b			Births attended by skilled health personnel ^c		Contraceptive prevalence rate ^d	
			Measles	DTP3	HepB3	At least 1 visit	At least 4 visits	Year	Year	Year	Year	
			(%) 2005	(%) 2005	(%) 2005	(%)	(%)	(%)				
136	Poland	EUR	98	99	98	100	2005
137	Portugal	EUR	93	93	94	100	2001
138	Qatar	EMR	99	97	97	62	58	1998	100	2002	43.2	1998
139	Republic of Korea	WPR	99	96	99	100	2003	80.5	1997
140	Republic of Moldova	EUR	97	98	99	99	...	1997	100	2005 ⁱ	62.4	2000
141	Romania	EUR	97	97	98	89	...	1999	98	2005	63.8	1999
142	Russian Federation	EUR	99	98	97	96	...	1999	99	2005	72.8	1999
143	Rwanda	AFR	89	95	95	93	10	2001	39	2005 ^k	13.2	2000
144	Saint Kitts and Nevis	AMR	99	99	99	...	100	2001	100	2005
145	Saint Lucia	AMR	94	95	95	...	100	2001	100	2004
146	Saint Vincent and the Grenadines	AMR	97	99	99	...	92	2001	100	2005
147	Samoa	WPR	57	64	60	100	2004
148	San Marino	EUR	94	95	95
149	Sao Tome and Principe	AFR	88	97	96	91	...	2000	79	2000	29.3	2000
150	Saudi Arabia	EMR	96	96	96	96	2004
151	Senegal	AFR	74	84	84	82	64	1999	52	2005	11.8	2005
152	Serbia	EUR	96	98	65
153	Seychelles	AFR	99	99	99
154	Sierra Leone	AFR	67	64	...	82	68	2001	42	2000	4.3	2000
155	Singapore	WPR	96	96	96	100	2004	62.0	1997
156	Slovakia	EUR	98	99	99	100	2004
157	Slovenia	EUR	94	96	100	2005
158	Solomon Islands	WPR	72	80	72	85	1999
159	Somalia	EMR	35	35	32	2001	25	2002
160	South Africa	AFR	82	94	94	89	72	1998	92	2003–2004	56.3	1998
161	Spain	EUR	97	96	96
162	Sri Lanka	SEAR	99	99	99	...	98	2001	97	2000	70.0	2000
163	Sudan	EMR	60	59	52	...	75	2001	57	1999 ^m
164	Suriname	AMR	91	83	83	91	91	2001	71	2000 ^k	42.1	2000
165	Swaziland	AFR	60	71	71	70	2000	27.7	2000
166	Sweden	EUR	94	99
167	Switzerland	EUR	82	93	100	2006
168	Syrian Arab Republic	EMR	98	99	99	...	51	2001	90	2004
169	Tajikistan	EUR	84	81	81	75	...	2000	71	2000	33.9	2000
170	Thailand	SEAR	96	98	96	...	86	2001	99	2000	72.2	1996–1997
171	The former Yugoslav Republic of Macedonia	EUR	96	97	53	99	2005
172	Timor-Leste	SEAR	48	55	18	2002	10.0	2003
173	Togo	AFR	70	82	...	78	46	1998	49	2000	25.7	2000
174	Tonga	WPR	99	99	99	98	2004
175	Trinidad and Tobago	AMR	93	95	95	96	98	2001	97	2002	38.2	2000
176	Tunisia	EMR	96	98	97	...	79	2001	90	2000	63.0	2001
177	Turkey	EUR	91	90	85	67	42	1998	83	2003	63.9	1998
178	Turkmenistan	EUR	99	99	99	87	83	2000	97	2000	61.8	2000
179	Tuvalu	WPR	62	93	79	100	2002
180	Uganda	AFR	86	84	84	92	40	2000–2001	39	2000–2001	22.8	2000–2001



Children aged <5 years sleeping under insecticide-treated bednets ^e		Antiretroviral therapy coverage		TB detection rate under DOTS ^h	TB treatment success under DOTS ⁱ	Children aged < 5 years with ARI symptoms taken to facility ^j		Children aged < 5 years with diarrhoea receiving ORT ^l		Children aged < 5 years with fever who received treatment with any antimalarial ^o		Children 6–59 months who received vitamin A supplementation ^l		Births by Caesarean section ^b	
(%)	Year	(%) Dec 2006	(%) Dec 2005			(%) 2005	(%) 2004 cohort	(%)	Year	(%)	Year	(%)	Year	(%)	Year
...		62	79	
...		85	84		30	2001
...		47	78		16	1998
...		18	80	
...		...	6	65	62	54.4	2005		6	1997
...		82	82		11	1999
...		11	84	30	59		12	1999
5	2000	72	36	29	77	26.9	2005	31.9	2005	12.6	2000	84.1	2005	2	2000
...		0	
...		92	64	
...		39	86	
...		66	100	
...		
23	2000		61.2	2000	
...		38	82	
2	2000	34	1	51	74	40.6	2005	52.5	2005	36.2	2000	75.3	2005	2	1997
...		79	91	
...		65	92	
2	2000	14	1	37	82		60.7	2000	...		2	1997
...		100	81	
...		39	88		18	2002
...		84	90		14	2002
...		55	87	
0	1999	<1	...	86	91		18.5	1999	
...		33	30	103	70	75.3	1998	89.1	1998		16	1998
...		
...		< 8	...	86	85	
0	2000	1	<1	35	77		50.2	2000	
3	2000	93	
0	2000	42	34	42	50		25.5	2000	
...		56	64		17	2001
...			10	2002
...		42	86	
2	2000	<5	12	22	84		68.9	2000	...		2	2002
...		88	...	73	74	
...		66	84		10	2001
4	2002	0	...	44	80		47.4	2002	
2	2000	24	8	18	67		60.0	2000	...		2	1998
...		96	
...		45	
...		82	90		8	2000
...		3	91	41.0	2003		14	1998
...		43	86		4	2000
...		35	100	
0	2000–2001	41	11	45	70	29.6	2001	53.1	2001	...		37.6	2001	3	2000–2001

Health service coverage

56835
83347-9473

Figures have been computed by WHO to ensure comparability; thus they are not necessarily the official statistics of Member States, which may use alternative rigorous methods.

	Member State	WHO region	Immunization coverage among 1-year-olds ^a			Antenatal care coverage ^b			Births attended by skilled health personnel ^c		Contraceptive prevalence rate ^d	
			Measles	DTP3	HepB3	At least 1 visit	At least 4 visits	Year	Year	Year	Year	
			(%) 2005	(%) 2005	(%) 2005	(%)	(%)					
181	Ukraine	EUR	96	96	97	90	...	1999	100	2005	67.5	1999
182	United Arab Emirates	EMR	92	94	92		100	2003	...	
183	United Kingdom	EUR	82	91		99	1998	84.0	2002
184	United Republic of Tanzania	AFR	91	90	90	96	69	1999	46	2004–2005	26.4	2004–2005
185	United States of America	AMR	93	96	92		99	2003	72.9	2002
186	Uruguay	AMR	95	96	96	...	94	2001	99	2004	...	
187	Uzbekistan	EUR	99	99	99		96	2000	67.7	2002
188	Vanuatu	WPR	70	66	56		87	2003	...	
189	Venezuela (Bolivarian Republic of)	AMR	76	87	88	...	90	2001	94	2000	...	
190	Viet Nam	WPR	95	95	94	70	29	2002	85	2002	78.5	2002
191	Yemen	EMR	76	86	86	34	11	1997	20	2003	20.8	1997
192	Zambia	AFR	84	80	80	94	71	2001–2002	43	2001–2002	34.2	2001–2002
193	Zimbabwe	AFR	85	90	90	82	64	1999	80	2005–2006	53.5	1999
194	The former state union of Serbia and Montenegro ^e	EUR		93	2001	58.3	2000

Region									
African Region	AFR	65	67	39		44	23.7
Region of the Americas	AMR	92	92	85		91	72.0
South-East Asia Region	SEAR	65	66	27		49	51.5
European Region	EUR	93	95	76		95	68.3
Eastern Mediterranean Region	EMR	82	82	74		53	39.9
Western Pacific Region	WPR	87	87	76		81	84.7
Global		77	78	55		63	61.9

... Data not available or not applicable; AFR, African Region; AMR, Region of the Americas; SEAR, South-East Asia Region; EUR, European Region; EMR, Eastern Mediterranean Region; WPR, Western Pacific Region.

The global values for rates and ratios are weighted averages; for absolute numbers they are the sums of all WHO regions.

^a DTP3, 3 doses of diphtheria–tetanus toxoid–pertussis vaccine; HepB3, 3 doses of Hepatitis B vaccine. *WHO/UNICEF estimates of national immunization coverage* [online database.]. Geneva, World Health Organization, 2006 (http://www.who.int/immunization_monitoring/routine/immunization_coverage/en/index4.html). Estimates based on data available up to August 2006. For countries recommending the first dose of measles among children older than 12 months of age, the indicator is calculated as the proportion of children less than 24 months of age receiving one dose of measles containing vaccine.

^b *The World Health Report 2005: make every mother and child count*. Geneva, World Health Organization, 2005 (<http://www.who.int/whr/2005/en/index.html>).

^c *WHO global database on births attended by skill health personnel*, Geneva, World Health Organization, 2007.

^d Percentage of women using contraception among those of reproductive age who are married or living with a partner. Source: *World contraceptive use 2005* [CD-ROM]. New York, Population Division, Department of Economic and Social Affairs, United Nations Secretariat, 2006. Additional updated information obtained by WHO's Department of Reproductive Health and Research directly from the UN Population Division.

^e *World malaria report 2005*. Geneva, World Health Organization, United Nations Children's Fund, 2005. Values for Cameroon and Chad have been updated for this report.

^f *Towards universal access: scaling up priority HIV/AIDS interventions in the health sector*. Geneva, World Health Organization, UNAIDS, United Nations Children's Fund, 2007. See Annex 1: Estimated number of people receiving antiretroviral therapy, people needing antiretroviral therapy and percentage coverage in WHO Member States. Ranges of estimates are available from this document.

^g PMTCT, preventing mother-to-child transmission. The coverage estimate is calculated by dividing the number of pregnant HIV-infected women who received antiretrovirals for PMTCT by the estimated unrounded number of pregnant HIV-infected women. Source: *Towards universal access: scaling up priority HIV/AIDS interventions in the health sector*. Geneva, World Health Organization, UNAIDS, United Nations Children's Fund, 2007. See Annex 3: Preventing mother-to-child transmission of HIV in low- and middle-income countries, 2005. Ranges of estimates are available from this document.



Children aged <5 years sleeping under insecticide-treated bednets ^e		Antiretroviral therapy coverage		TB detection rate under DOTS ^h	TB treatment success under DOTS ⁱ	Children aged < 5 years with ARI symptoms taken to facility ^j		Children aged < 5 years with diarrhoea receiving ORT ^j		Children aged < 5 years with fever who received treatment with any antimalarial ^e		Children 6–59 months who received vitamin A supplementation ^j		Births by Caesarean section ^b	
(%)	Year	(%) Dec 2006	(%) Dec 2005			(%) 2005	(%) 2004 cohort	(%)	Year	(%)	Year	(%)	Year	(%)	Year
...		...	90		9	1999
...		19	70	
...			17	1997
2	1999	18	6	45	81	56.6	2005	70.0	2005	53.4	1999	41.2	2005	3	1999
...		85	61		23	2000
...		51	...	83	
...		39	78	
...		61	90	
...		71	...	73	81	
16	2000	17	7	84	93	71.3	2002	74.0	2002	6.5	2000	...		10	2002
...		0	...	41	82	32.2	1997	64.6	1997		1	1997
7	2001–2002	35	15	52	83	69.1	2002	66.9	2002	51.9	2001–2002	67.4	2002	2	2001–2002
...		15	9	41	54	49.8	1999	79.7	1999		7	1999
...		

...	29	11	50	74	
...	65	80	
...	18	6	64	87	
...	35	74	
...	44	83	
...	30	4	76	91	
...	60	84	

^h TB, tuberculosis; DOTS, internationally recommended TB control strategy. The detection rate is the number of new smear-positive cases notified to WHO divided by the estimated number of new smear-positive cases. Source: *Global tuberculosis control: surveillance, planning, financing. WHO report 2007*. Geneva, World Health Organization, 2007 (WHO/HTM/TB/2007.376) (http://www.who.int/tb/publications/global_report).

ⁱ The treatment success rate is the percentage of new smear-positive patients registered for treatment under DOTS during 2004 who were cured (with laboratory confirmation) or completed their course of treatment. Source: *Global tuberculosis control: surveillance, planning, financing. WHO report 2007*. Geneva, World Health Organization, 2007 (WHO/HTM/TB/2007.376) (http://www.who.int/tb/publications/global_report).

^j ARI, acute respiratory infection; ORT, oral rehydration therapy. Data compiled by the Department of Child and Adolescent Health and Development, WHO, from *Demographic and Health Surveys: final reports, 2007* (http://www.measuredhs.com/pubs/search/search_results.cfm?Type=5&srchTp=type&newSrch=1, accessed 15 February 2007).

^k In this case, data do not relate to “skilled health personnel” as defined in the document *Making pregnancy safer: the critical role of the skilled attendant. A joint statement by WHO, ICM and FIGO*. Geneva, World Health Organization, 2004. Further information can be found at http://www.who.int/reproductive-health/global_monitoring/RHRxmls/RHRmainpage.htm.

^l Excludes Transnistria region.

^m Covers Northern Sudan and selected sites in Southern Sudan.

ⁿ See footnote o to the table on Health status: mortality.